

## RESEARCH ARTICLE

# Diversity of putative ericoid mycorrhizal fungi increases with soil age and progressive phosphorus limitation across a 4.1-million-year chronosequence

Devin R. Leopold<sup>1,2,\*†</sup>, Kabir G. Peay<sup>1</sup>, Peter M. Vitousek<sup>1</sup> and Tadashi Fukami<sup>1</sup>

<sup>1</sup>Department of Biology, Stanford University, 371 Serra Mall, Stanford, CA 94305–5020, USA and <sup>2</sup>Department of Botany and Plant Pathology, Oregon State University, 4575 SW Research Way, Corvallis, OR 97331, USA

\*Corresponding author: OSU Botany and Plant Pathology, 4575 SW Research Way, Corvallis, OR 97333, USA. Tel: 603-702-1203; E-mail: [devin.leopold@gmail.com](mailto:devin.leopold@gmail.com)

**One sentence summary:** The diversity and species composition of fungal symbionts associated with the roots of ericaceous plants depends on soil age and nutrient availability.

Editor: Ian Anderson

<sup>†</sup>Devin R. Leopold, <http://orcid.org/0000-0003-2124-2116>

## ABSTRACT

Ericaceous plants rely on ericoid mycorrhizal fungi for nutrient acquisition. However, the factors that affect the composition and structure of fungal communities associated with the roots of ericaceous plants remain largely unknown. Here, we use a 4.1-million-year (myr) soil chronosequence in Hawaii to test the hypothesis that changes in nutrient availability with soil age determine the diversity and species composition of fungi associated with ericoid roots. We sampled roots of a native Hawaiian plant, *Vaccinium calycinum*, and used DNA metabarcoding to quantify changes in fungal diversity and community composition. We also used a fertilization experiment at the youngest and oldest sites to assess the importance of nutrient limitation. We found an increase in diversity and a clear pattern of species turnover across the chronosequence, driven largely by putative ericoid mycorrhizal fungi. Fertilization with nitrogen at the youngest site and phosphorus at the oldest site reduced fungal diversity, suggesting a direct role of nutrient limitation. Our results also reveal the presence of novel fungal species associated with Hawaiian Ericaceae and suggest a greater importance of phosphorus availability for communities of ericoid mycorrhizal fungi than is generally assumed.

**Keywords:** chronosequence; ericoid mycorrhizae; mycorrhizal diversity; nutrient limitation; pedogenesis; root-associated fungi

## INTRODUCTION

The plant family Ericaceae includes clades that are largely defined by morphologically and functionally distinct mycorrhizal symbioses, with the largest group forming a unique endomycorrhizal association known as an ericoid mycorrhiza (ErM; Pearson and Read 1973a; Perotto et al. 2012; Leopold 2016). The ErM symbiosis allows ericaceous plants to proliferate in

harsh environments where acidic soils, low temperatures, or excessive soil moisture slow the degradation of organic matter and limit mineral nutrient availability (Read 1991; Cairney and Meharg 2003; Mitchell and Gibson 2006). In these habitats, which can be found on every continent except Antarctica (Kohout 2017), ericoid mycorrhizal fungi (ErMF) facilitate nutrient cycling by degrading complex organic matter and providing

Received: 27 August 2020; Accepted: 27 January 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of FEMS. All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

their hosts with access to nutrient pools that would be unavailable otherwise (Wurzburger, Higgins and Hendrick 2012; Adamczyk et al. 2016; Perotto, Daghino and Martino 2018). Ericaceous plants and ErMF also contribute to carbon sequestration in these environments through the production of recalcitrant plant litter and hyphal necromass (Clemmensen et al. 2013, 2015). However, despite the ecological importance of the ErM symbiosis, and significant functional variation within and among ErMF species (Cairney et al. 2000; Whittaker and Cairney 2001; Grelet et al. 2009b; Wurzburger, Higgins and Hendrick 2012), little is known about how factors such as soil nutrient availability affect the community composition of ErMF and other fungi associated with ericoid roots.

Currently, most data on ErMF originates from temperate and boreal regions, where cooler temperatures and repeated glaciation throughout the Pleistocene have maintained widespread nitrogen (N) limitation, even in late-successional ecosystems (Tamm 1991; Vitousek and Howarth 1991). As a result, research on ErMF has focused on their ability to utilize various organic N sources and influence host N status (Leake and Read 1991; Nasholm et al. 1998; Xiao and Berch 1999; Cairney et al. 2000; Grelet et al. 2009b; Ishida and Nordin 2010). However, the distribution of ericaceous plants is not limited by these habitat conditions (Leopold 2016; Kohout 2017) and there is evidence that ErMF also facilitate the uptake of phosphorus (P) and other mineral nutrients for host plants (Pearson and Read 1973b; Read 1983; Myers and Leake 1996). This observation suggests that the relative availability of limiting nutrients in soil, P as well as N, may influence ericaceous plant-fungal interactions and ErMF community composition (Hazard et al. 2014; Van Geel et al. 2020).

One source of natural variation in the availability of N and P in soil is pedogenesis, or long-term soil development. Over hundreds of thousands to millions of years, and in the absence of rejuvenating disturbance (Peltzer et al. 2010), pedogenesis causes predictable changes in ecosystem-level nutrient limitation, primarily through the accumulation of N from biological inputs and the progressive loss and occlusion of rock-derived nutrients, especially P (Walker and Syers 1976; Chadwick et al. 1999). These changes result in an initial, progressive phase, defined by N limitation of primary productivity and the rapid accumulation of organic matter; a mature phase, defined by maximal productivity and co-limitation by N and P; and a retrogressive phase, defined by P limitation and reduced rates of nutrient cycling (Vitousek and Farrington 1997; Richardson et al. 2004; Wardle 2004; Peltzer et al. 2010).

Because the direct observation of long-term soil development in a single site is not possible, soil chronosequences, or gradients of soil age in which other putative controlling factors (i.e. parent material, climate, vegetation type, etc.) are held reasonably constant, can provide insight into the effects of changes in nutrient availability with soil age (Fukami and Wardle 2005; Lambers et al. 2008; Walker et al. 2010; Dickie et al. 2013). Soil chronosequence studies of arbuscular mycorrhizal fungi (AMF) have suggested direct N and P limitation of fungi in young and old ecosystems, respectively (Treseder and Allen 2002), and a peak in AMF richness at fertile, middle-aged sites (Krüger et al. 2015). However, AMF are primarily involved in mineral nutrient uptake (Smith and Read 2006) and appear to lack the extracellular enzymes needed to access the organic nutrient pools available to some ErMF (Tisserant et al. 2013). ErMF diversity could respond differently to changing mineral nutrient availability with soil aging, but the relative importance of N and P limitation for ErMF is currently unknown. Furthermore, despite adaptation of the ErM symbiosis to soils with low mineral nutrient

availability, studies of ErMF from retrogressive, P-limited habitats are lacking (Dickie et al. 2013).


To explore the effects of long-term soil development on ErMF, and the broader community of ericaceous root-associated fungi, we sampled roots of a single ericaceous plant species, *Vaccinium calycinum*, across a 4.1-million-year (myr) soil chronosequence in the Hawaiian Islands, known as the Long Substrate Age Gradient (LSAG; Vitousek 2004). The LSAG is a useful system to study ErMF for three reasons. First, *V. calycinum*, a common host plant species, is present at all LSAG sites. The prevalence of a common host species across the chronosequence controls for changes in host identity, which can influence the composition of mycorrhizal symbiont communities (Martínez-García et al. 2015), though intraspecific variation in host traits affecting plant-fungal interactions are also possible (Johnson et al. 2010). Second, vegetation at LSAG sites is dominated by a single canopy tree species and a common suite of understory plants (Kitayama and Mueller-Dombois 1995), which all form mycorrhizal associations with AMF (Koske, Gemma and Flynn 1992). The only other ericaceous species present at any of the LSAG sites, *Vaccinium dentatum* and *Leptocophylla tameiameiae*, occur very sparsely at all sites, which further limits the potential impact of changes in host vegetation along the chronosequence. Third, changes in nutrient limitation across the LSAG have been experimentally demonstrated through N and P fertilizer addition experiments (Vitousek 2004). Because these fertilization experiments are long-term (>20 years) and ongoing at the oldest and youngest LSAG sites, this system presents a unique opportunity to experimentally test whether ecosystem-level nutrient limitation (i.e. N and P availability in young and old sites, respectively) plays a role in structuring communities of ericaceous root-associated fungi.

We began with the hypothesis that the diversity of fungi associated with ericaceous roots increases throughout long-term soil development, with ErMF diversity increasing owing to the accumulation of complex organic nutrient pools. We predicted that diversity would increase most rapidly during the early stages of pedogenesis due to the formation and development of an organic soil horizon and that an increase would continue into the retrogressive stages as soil weathering and occlusion of mineral nutrients promotes niche partitioning in the rhizosphere (Turner 2008; DeForest and Scott 2010). We also used long-term fertilization experiments to test the hypothesis that nutrient limitation of primary productivity, N at the youngest site and P at the oldest site, shapes the species composition of ErMF and other ericaceous root-associated fungi.

## MATERIALS AND METHODS

### Study system

The LSAG chronosequence consists of six study sites that were established to address questions about the mechanisms that control nutrient cycling and limitation throughout long-term soil development by exploiting the increasing age of Hawaiian volcanoes with increasing distance from the active hot spot (Vitousek 2004). Samples for the study reported here were collected from five LSAG sites ranging in substrate age from 300 year (yr) to 4.1 myr (Fig. 1). One site (1.4 myr) was not sampled due to logistical constraints. In this system, substrate age is an estimate of how long ago the parent substrate was deposited by volcanic activity. The locations of the LSAG sites were chosen to minimize variation in state factors other than the age of the soil substrate. All sites share similar initial parent substrate, occur



**Figure 1.** Map of the main Hawaiian Islands showing the locations and ages of the five sites in the Long Substrate Age Gradient chronosequence used in the current study.

on the constructional slope of a shield volcano, are ca. 1200 m asl, receive ca. 2500 mm rain annually and have a mean annual temperature of ca. 16°C (for detailed site descriptions, see Crews et al. 1995; Vitousek 2004). Biotic variation is also highly constrained across the LSAG. Vegetation consists of native forest dominated by the canopy tree species *Metrosideros polymorpha* (Myrtaceae), with an understory that includes *Cheirodendron trigynum*, *Cibotium* spp. (tree ferns), *Coprosma* spp., *Ilex anomala*, *Myrsine* spp. and *V. calycinum* (Kitayama and Mueller-Dombois 1995).

In addition to the observational data collected across the LSAG chronosequence, fertilizer addition experiments were established in some LSAG sites to experimentally test hypotheses concerning the dynamics of nutrient limitation to ecosystem processes over geologic time scales. Fertilization plots sampled for the current study were established at the 300-yr site in 1985 (Vitousek et al. 1993) and at the 4.1-my site in 1991 (Herbert and Fownes 1995), and so had been ongoing for 30 and 23 years, respectively, at the time of this sampling. At each site, 16 plots (15 × 15 m) were randomly assigned to either an unfertilized control or treatments that received fertilizer in the form of N (initially half ammonium nitrate and half urea, since 2002 just urea), P (triple superphosphate), or both N and P. Fertilizer was applied twice a year until 2002 and biannually thereafter at a consistent annual rate of 100 kg/ha/yr N and 100 kg/ha/yr P.

### Sample collection

Sampling occurred in August 2014, with the exception of the nutrient addition plots at the 300-yr site, which were sampled in August 2013 as part of a preliminary feasibility study. To account for possible inter-annual variation, an additional 12 plants from non-fertilized plots at the 300-yr site were collected in 2014. Initial analyses suggested that samples from unfertilized plots collected at the 300-yr site in different years had similar fungal communities. Nonetheless, comparisons across the chronosequence reported here utilize only the 2014 samples and analy-

ses of samples from fertilizer plots involved comparisons within each site, using the 2013 samples for the 300-yr site and the 2014 samples for the 4.1-my site.

At each LSAG site, 12 mature *V. calycinum* plants were randomly selected within a 200 × 200 m area. In the fertilizer addition plots at the 300-yr and 4.1-my sites, two plants were sampled in each 15 × 15 m plot, resulting in 32 samples per site (eight samples per fertilizer treatment in each site). Plant were selected to be similar in size (ca. 0.5-1 m tall) and for each plant a portion of the root system (ca. 25 × 25 × 10 cm) and the adhering soil was removed with a hand trowel and bagged for transport. Samples were refrigerated (4°C) until they could be processed, always within 48 h of collection. Fine terminal roots were manually separated from the soil and rinsed in tap water to remove all visible soil particles and 12 segments, ca. 2 cm each, were randomly selected and pooled for each plant. Pooled root samples were surface sterilized by sequential vortexing for 1 min in sterile water, 70% EtOH, 50% household bleach and then rinsed three times in sterile water. Surface sterilized roots were then stored at -80°C.

### Fungal community metabarcoding

Frozen roots were homogenized by bead beating in CTAB lysis buffer and total DNA was extracted using the Nucleospin Plant II kit (Macherey-Nagel, Bethlehem, PA, USA). In order to provide negative controls for Illumina sequencing, four sterile bead beating tubes were processed in parallel with the tubes containing root samples. Following DNA extraction, the first half of the internal transcribed spacer region (ITS1) of fungal nrRNA was amplified using Illumina fusion PCR primers. Primers included the ITS1 primers ITS1F (forward) or ITS2 (reverse), modified to include Illumina adapters and a sample specific 12 bp, error-correcting Golay barcode (Smith and Peay 2014). PCR was carried out in 25 µL reactions using 1 µL of template DNA (diluted 1:20), 0.5 µL of each 10 µM primer, 12.5 µL OneTaq Hot-Start 2X Master Mix (New England Biolabs, Ipswich, MA, USA) and a cycling pro-

gram consisting of: initial denaturing at 94°C (1 min), 30 cycles of 95°C (30 s), 52°C (30 s) and 68°C (30 s) and a final elongation stage of 68°C (5 min). PCR reactions were carried out in triplicate for each sample and were individually checked for successful amplification using gel electrophoresis. The individual PCR reactions for each sample were cleaned and normalized to 2.5 ng/μL using a Just-a-Plate, 96 well normalization and purification plate (Charm Biotech, San Diego, CA, USA). Samples were then pooled and sequenced using Illumina MiSeq paired-end sequencing (2 × 300 bp) at the Stanford Functional Genomics Facility. Raw sequence data were deposited in the NCBI short read archive under BioProject ID PRJNA548137.

## Bioinformatics

Preliminary analyses suggested that the reverse read quality was poor, so only the forward reads were used for all analyses. Raw reads were first trimmed to remove 3' gene primer and sequencing adapter contamination using cutadapt (Martin 2011). The trimmed reads were then pooled and denoized using DADA2 (Callahan et al. 2016), excluding reads with an expected error rate >2 bp, a length <75 bp and putative chimeric sequences. Denoized reads were then collapsed to 99% operational taxonomic units (OTUs) using agglomerative, single-linkage clustering implemented in DECIPHER (Wright 2016). The OTU table was then filtered of likely contaminants by removing any OTU with an average abundance across all samples that was less than the maximum abundance in any of the negative controls.

Taxonomic predictions for OTUs were made using the DADA2 implementation of the RDP classifier (Wang et al. 2007) trained on the UNITE v8.2 species hypothesis database (Abarenkov et al. 2020). Because a prominent group of ErMF, members of the *Hyaloscypha hepaticicola* aggregate (= *Hymenoscyphus ericae* aggregate; = *Rhizoscyphus ericae* aggregate), has undergone numerous taxonomic revisions (Vrålstad, Fosshem and Schumacher 2000; Vrålstad, Myhre and Schumacher 2002; Hambleton and Sigler 2005), we manually curated the relevant taxonomy in the training data. We resolved inconsistencies in this clade following the most recent updates reported by Fehrer et al. (2019), which consolidated members of the *H. hepaticicola* aggregate in the genus *Hyaloscypha*.

Identification of potential ErMF in the OTU database was complicated by two factors. First, the short read length and high variability of the ITS region, combined with limited representation of both Hawaiian root/soil associated fungi and fungi associated with ericaceous plants in current reference databases resulted in low confidence for fine scale taxonomic assignments. Second, there is significant uncertainty about the taxonomic range of fungal species capable of forming ErM. We chose an inclusive approach, defining the subset of putative ErMF to include all OTUs assigned to fungal orders reported to include ericoid fungi (Leopold 2016). An alternative approach, manually curating putative ErMF OTUs by looking at the sources of the individual sequences comprising the closest matching UNITE species hypotheses yielded qualitatively similar results. We focus here on the approach using fungal orders as it is more easily replicated. However, either approach is likely to include many non-mycorrhizal endophytes, which may have overlapping functional attributes with ErMF (Newsham 2011), as well as saprobes, and parasites, and it is not currently possible to definitively distinguish ErMF taxa with sequence data alone.

## Statistical analysis of chronosequence data

To assess if the fungal diversity changed across the LSAG, we estimated alpha diversity at the sample level for both the complete data set and the subset of putative ErMF. To account for unequal sequencing depth among samples we estimated alpha diversity using Hill numbers, or the effective number of species, at a sampling depth of 5000 sequences per sample using the R-package iNEXT (Chao et al. 2014; Hsieh, Ma and Chao 2016). We calculated Hill numbers for each sample using a scaling of 0 (equivalent to species richness) and a scaling of 1 (equivalent to the exponent of Shannon entropy, or Shannon diversity) and analyzed the results in parallel. Variation in alpha diversity among chronosequence sites was tested using ANOVAs and *F*-tests, followed by Tukey's HSD post-hoc tests.

To determine how the composition of fungal communities changed across the LSAG, we first calculated Bray–Curtis dissimilarity among samples using the proportional abundance of OTUs in each sample to account for unequal sampling depth (McMurdie and Holmes 2014). We initially tested alternative dissimilarity metrics, including the binary Jaccard distance, the probabilistic Raup–Crick distance and the information-theoretic based Jensen–Shannon Divergence, and found that all produced qualitatively similar results to those obtained with Bray–Curtis dissimilarity, which we will focus on here. Dissimilarities were visualized using non-metric multidimensional scaling (NMDS) implemented in the R-package *vegan* (Oksanen et al. 2017). Contours representing increasing site age were fit to the 2-dimensional NMDS ordination, using the *vegan* function *ordisurf*, to aid interpretation of species turnover across the chronosequence. The significance of changes in community composition were tested with a permutational multivariate analysis of variance (Anderson 2001). Both  $\log_{10}$ -transformed site age and site identity as a factor were used as predictors in perMANOVAs to assess both the significance of variation in community composition among sites and the proportion of this variation that could be explained by turnover along the chronosequence. OTUs occurring at individual LSAG sites with greater abundance and more often than expected by chance alone were identified using indicator species analysis (Dufrene and Legendre 1997) using the R-package *labdsv* (Roberts 2019), and visualized on a bipartite graph of the 50 most abundant OTUs.

## Statistical analysis of fertilizer plot data

To test the effects of fertilization on fungal composition and diversity, we analyzed both the complete data set and the subset of putative ErMF OTUs from the fertilizer addition plots at the 300-yr and 4.1-myr LSAG sites. We followed the procedure described above to estimate fungal richness and Shannon diversity and then used ANOVAs and *F*-tests to assess the direct and interactive effects of N and P addition. Although fertilizer treatments were applied to 16 spatially randomized plots at each site, with the original goal of addressing ecosystem-level responses to nutrient addition (see, Vitousek 2004 and references therein), we focused our analyses on individual plants ( $n = 32$ ) as the experimental unit of interest in this study. This choice was justified because *V. calycinum* is sparsely distributed in the understory and the plants targeted for sampling were always physically separated (>5 m) within plots, without a contiguous network of ErMF hosts. Furthermore, preliminary analysis indicated a lack of within-plot correlation of fungal richness or

diversity. To test whether community composition was affected by fertilization, we used NMDS and permutational MANOVAs of Bray–Curtis dissimilarity calculated on proportional OTU abundance.

## Reproducibility of analyses

All statistical analyses were conducted in the R v4.2 statistical computing environment (R Core Team 2020), using the package *tidyverse* (Wickham et al. 2019) for general data manipulation and *phyloseq* (McMurdie and Holmes 2013) to manipulate OTU tables and associated metadata. Code for reproducing all bioinformatic processing, statistical analyses and figures, along with all necessary sample metadata, has been publicly archived (DOI: 10.5281/zenodo.3979769).

## RESULTS

### Identification of fungal OTUs

From 4.6 M raw reads, we retained 2.2 M high-quality sequences after bioinformatic processing, from which we identified 685 fungal OTUs. The subset of putative ErMF included 30.8% of the total OTUs, and accounted for 66.5% of all sequences. Many taxa could not be confidently assigned to genera or family (Fig. 2), and the lack of high similarity matches for many OTUs in the UNITE database suggests the presence of many novel taxa. For example, the most abundant taxon we identified (OTU.1) shares only 85% sequence similarity with the nearest sequence in the UNITE database and could not be confidently assigned beyond the order Helotiales. The second most abundant taxon (OTU.2) was assigned to the order Trechisporales and shares 99% sequence similarity with a UNITE species hypothesis that contains only a few individuals sequenced without culturing from the roots of plants in the Orchidaceae and Clustaceae, originating from Eastern Africa and Taiwan. However, the next closest UNITE species hypothesis in the Trechisporales shares only 75% sequence similarity, suggesting that this is could be a specious taxonomic assignment (see discussion, below). We did identify numerous taxa assigned to the *H. hepaticicola* aggregate based on their ITS1 sequences, including OTUs classified as the archetypal ErMF, *H. hepaticicola* ( $\equiv$  *Rhizoscyphus ericae* &  $\equiv$  *Pezoloma ericae*; Fehrer et al. 2019), though this taxon was notably absent at the youngest site (Fig. 2). The common ErMF species, *Oidiendron maius*, was only observed in very low abundance in the sequence data, primarily in samples from the oldest chronosequence site. No members of the Sebaciniales were observed, despite their common occurrence as root endophytes and ErMF on ericaceous plants (Selosse et al. 2007; Vohník et al. 2016).

### Site age effects on fungal diversity and composition

Fungal OTU richness and diversity were greatest in *V. calycinum* roots collected from the oldest chronosequence site (Fig. 3). When all fungi were considered together (Fig. 3A), variation in OTU richness ( $F_{4,55} = 5.14$ ;  $P = 0.001$ ) was primarily the result of lower richness in the more fertile, middle-aged sites, while variation in Shannon diversity ( $F_{4,55} = 11.9$ ;  $P < 0.001$ ) was primarily due to greater richness at the oldest site. For putative ErMF (Fig. 3B), variation in OTU richness ( $F_{4,54} = 5.62$ ;  $P < 0.001$ ) was the result of an increase at the two oldest sites, relative to the younger sites, while variation in Shannon diversity ( $F_{4,54} = 11.4$ ;  $P < 0.001$ ) was the result of a progressive increase with site age.

Fungal communities associated with *V. calycinum* roots differed in composition among chronosequence sites for all fungi ( $F_{4,55} = 6.92$ ,  $r^2 = 0.33$ ,  $P < 0.001$ ) and the subset of putative ErMF ( $F_{4,55} = 9.42$ ,  $r^2 = 0.40$ ,  $P < 0.001$ ). Similarly aged sites tended to have more similar OTU compositions (Fig. 4), and site age explained roughly a third of the variation in OTU composition that was attributed to variation among sites, both for all fungi ( $F_{1,58} = 7.64$ ,  $r^2 = 0.12$ ,  $P < 0.001$ ) and putative ErMF ( $F_{1,58} = 9.41$ ,  $r^2 = 0.14$ ,  $P < 0.001$ ). In addition, while many of the most abundant OTUs were broadly distributed across the chronosequence, occurring in most or all sites, distribution patterns varied among taxa and many could be identified as being significantly associated with individual LSAG sites (Fig. 2).

### Fertilization effects on fungal composition and diversity


The effect of fertilization on fungal richness and diversity depended on site (Fig. 5; Tables S1 and S2, Supporting Information). At the 300-yr-old site, the addition of N reduced both richness and Shannon diversity for all fungi ( $F_{1,28} = 10.1$ ;  $P = 0.004$  &  $F_{1,28} = 5.95$ ;  $P = 0.02$ , respectively). For putative ErMF, the addition of N at the 300-yr-old site also reduced OTU richness ( $F_{1,28} = 12.9$ ;  $P = 0.001$ ), and marginally reduced Shannon diversity ( $F_{1,28} = 3.37$ ;  $P = 0.08$ ). In contrast, at the 4.1-myrr site it was the addition of P that reduced OTU richness and Shannon diversity for all fungi ( $F_{1,28} = 6.65$ ;  $P = 0.02$  &  $F_{1,28} = 15.8$ ;  $P < 0.001$ , respectively) and putative ErMF ( $F_{1,28} = 3.60$ ;  $P = 0.07$  &  $F_{1,28} = 8.7$ ;  $P = 0.006$ , respectively).

In contrast to richness and diversity, fungal community composition was not primarily impacted by addition of the limiting nutrient at each site (Table 1; Figure S1, Supporting Information). Instead, both N and P additions affected fungal composition, but P addition had the greatest impact on composition at both sites, for all fungi and the subset of putative ErMF.


## DISCUSSION

Consistent with our initial hypothesis, we found that *V. calycinum* associates with more putative ErMF species in sites with older soils. This pattern was not simply the result of increasing richness, but also greater evenness, which could reflect increased niche partitioning in response to the accumulation of soil organic matter and the development of a structured organic soil horizon during pedogenesis (Goh et al. 1976; Torn et al. 1997). Although speculative, this explanation is consistent with the fact that ErMF taxa can vary in their affinity for specific nutrient sources (Cairney et al. 2000; Whittaker and Cairney 2001; Grelet, Meharg and Alexander 2005; Grelet et al. 2009b) and can be spatially structured across soil layers (Wurzburger, Higgins and Hendrick 2012). However, it is also possible that dispersal limitation (Cline and Zak 2014), *in-situ* diversification (Gillespie 2016) and enhanced competitive equivalence (Leopold and Fukami 2020) contributed to the greater diversity observed in older sites. Manipulative experiments would be needed to quantify the role of niche partitioning relative to other factors.

Results from our fertilization experiment suggest that soil-age related nutrient limitation influenced fungal richness and diversity. Specifically, we found that richness and diversity decreased in response to the addition of N and P at the youngest and oldest chronosequence sites, respectively. Previous studies have shown that plant growth and net primary productivity are limited by N at the 300-yr site and P at the 4.1-myrr site (Harrington, Fownes and Vitousek 2001; Vitousek 2004), and P limits rates of litter decomposition and soil organic matter processing



**Figure 2.** Bipartite network showing the relative abundances of the 50 most common fungal OTUs associated with the roots of *Vaccinium calycinum* across the Long Substrate Age Gradient. The widths of connections between sites (left) and OTUs (right) shows the mean relative abundance of the OTU at that site and the size of nodes representing OTUs is proportional to the sum of its mean relative abundances across all sites. Relative abundances were calculated using only OTUs depicted in this figure, such that site nodes do not vary in size. Colored OTU nodes and connections identify significant site associations (indicator species analysis). Taxonomic predictions for each OTU are indicated following the unique OTU identifier.



**Figure 3.** Variation in species richness and Shannon diversity of (A) all fungi and (B) putative ErMF associated with *Vaccinium calycinum* roots across the Long Substrate Age Gradient. Boxplots show the distribution of the point estimates of richness and diversity for 12 replicate samples at each site. Points indicate the predicted mean ( $\pm$  95% bootstrapped CI) species richness or diversity. Letters indicate significant pairwise differences in post-hoc comparisons.


in the 4.1-myr site (Hobbie and Vitousek 2000; Reed, Vitousek and Cleveland 2011). Moreover, Treseder and Allen (2002) found that growth of arbuscular mycorrhizal fungi associating with non-ericaceous plants in this system is directly N-limited at the 300-yr site and P-limited at the 4.1-myr site. Our results could suggest that ErMF, in contrast with arbuscular mycorrhizal fungi, are not directly limited by N and P availability in the LSAG, possibly due to their ability to access recalcitrant organic nutrient pools (Read 1983, 1991; Perotto, Daghino and Martino 2018). Instead, changes in plant growth with fertilizer addition may change the relative strength of carbon-sinks above- and below-ground, causing the host plant to reallocate resources (i.e. carbon-rich photosynthates) away from root-associated symbionts (Olsson, Rahm and Aliasgharizad 2010; Kiers et al. 2011; Konvalinková et al. 2017), resulting in reduced fungal richness and diversity in the rhizosphere.

Alternative explanations for the effects of fertilization on ErMF richness and diversity are also possible. For example, if niche partitioning was important for ErMF community structuring, fertilization could reduce fungal diversity by skewing the available nutrient pools towards mineral sources, or by altering root morphology or depth. It is also possible that competition with other rhizosphere or soil-associated fungi contributed to changes in ErMF richness and diversity. Koske et al. (1990) reported that *V. calycinum* and other native Hawaiian ericoid plants could host both ErM and arbuscular mycorrhizae, suggesting competition for host resources. Finally, changes in plant community composition with fertilization could have affected the physical and chemical composition of the soil environment, as well as the local pool of rhizosphere-associated microbiota. Ostertag and Verville (2002) reported that the relative abundance of the invasive plants *Hedychium gardnerianum* and *Rubus argutus* increased with fertilization at the 300-yr and 4.1-myr sites, respectively. However, the composition of the native plant community, including the relative abundance of *V. calycinum*, was largely unaffected by fertilization. Regardless of the mechanism,

reduced ErMF diversity under fertilizer addition could have significant implications for stability of ERM symbioses in the context of anthropogenic nutrient deposition in the nutrient-poor habitats where ericaceous plant often thrive (Van Geel et al. 2020).

Although we observed a clear pattern of turnover in species composition across the soil chronosequence (Fig. 4), results from our fertilization experiment suggest that host and ecosystem-level nutrient limitation are not the primary determinants of fungal species composition in this system (Table 1). This could be because dispersal limitation restricts the pools of ErMF species present at each site (Hutton et al. 1997), limiting the effect of localized nutrient additions. Nonetheless, both N and P additions did influence fungal community composition at both sites, though the effects of P additions were consistently larger, even at the N-limited site. Previous work in this system has shown that fertilization with P at the 4.1-myr site significantly affects litter chemistry, whereas fertilization with N at the 300-yr site does not (Harrington, Fownes and Vitousek 2001). Because ErMF may be directly accessing recalcitrant nutrients in the upper, organic soil layers (Read, Leake and Perez-Moreno 2004), changes in litter chemistry may explain why the strongest fertilization effects we observed for putative ErMF with P addition at the 4.1-myr site. These results highlight the relative lack of data on ErM in P-limited, tropical systems and suggest that P availability may play a greater role in determining the abundance and distribution of ErMF than is generally assumed.

We do not have the histological data to determine the mycorrhizal status of the fungal OTUs we identified. However, our results suggest that many novel ErMF species could occur in this system. For example, the two most abundant OTUs in our study, which accounted for 32% of all reads across the chronosequence and 64% of the reads at the youngest site (Fig. 2), were not closely related to any well-described ErMF species. Although the most abundant OTU shared only 85% sequence similarity with the nearest taxon in the UNITE database, which comprises a large




**Figure 4.** Non-metric multidimensional scaling of community dissimilarity (Jensen Shannon distance) for (A) all fungi (stress = 0.26) and (B) putative ErMF (stress = 0.25) associated with *Vaccinium calycinum* roots collected across a 4.1-myrr chronosequence. Points represent individual plants, with color indicating the age of the site where the sample was collected. Contours indicate a smooth surface of  $\log_{10}$ -transformed site age fit to the ordination space, where increasing site age is indicated by darker contours.

group of poorly classified Helotiales, many of the most similar taxa were isolated from the roots of ErM or ecto-mycorrhizal plants, which can host ErMF as endophytes or mycorrhizal fungi capable of forming both ecto- and ericoid mycorrhizae (Villarreal-Ruiz, Anderson and Alexander 2004; Vrålstad 2004; Grelet et al. 2009a; Villarreal-Ruiz et al. 2012; Vohník et al. 2013). There also is considerable reason to doubt the taxonomic prediction for the second most abundant OTU as Trechisporales. A mycorrhizal symbiont forming a unique, sheathed-ericoid mycorrhizal symbiosis in Norway was also originally thought to be most closely related to the Trechisporales (Vohník et al. 2012). This taxon was later identified as *Kurtia argillacea*, in the Hymenochaetales, based on multi-gene phylogenetic analyses, which showed that proper taxonomic placement using nrRNA genes was not possible for this taxon (Kolařík and Vohník 2018). Although OTU.2 in our study does not share substantive sequence similarity with the one available ITS sequence for *Kurtia argillacea*, it does share some characteristics, including an elevated GC content (57%), which Kolařík and Vohník (2018) suggest could be the result of a non-standard substitution rate, possibly

related to the transition to a mycorrhizal lifestyle. Culture-based investigations of this taxon and other novel taxa associated with Hawaiian ericaceous plants are needed to definitively address the possibility that our data set includes novel ErMF. Although it is also possible that some of these species are non-mycorrhizal endophytes, this possibility does not preclude a role in nutrient uptake or other beneficial functions for the host plant.

In addition to highlighting the potential for novel ErMF taxa colonizing Hawaiian Ericaceae, our study also provides new data on the distributions of some common ErMF. For example, we identified the prominent ErMF species, *H. hepaticicola*, extending the known range of this species (Bruzzone et al. 2017). Although *H. hepaticicola* was observed in four chronosequence sites, it was not identified at the youngest site, which may be the result of dispersal limitation or environmental filtering. In contrast, we did not detect the presence of any fungi in the order Sebaciales, which includes some widespread ErMF, as well as non-mycorrhizal endophytes commonly associated with ericoid roots (Selosse et al. 2007; Vohník et al. 2016), though we cannot rule out the possibility that our choice of PCR primers introduced



**Figure 5.** The effect of fertilization with nitrogen (N), phosphorus (P), or both (NP) on the species richness and Shannon diversity of fungal OTUs associated with the roots of *Vaccinium calycinum* at (A) the 300-yr and (B) the 4.1-myrr sites in the Long Substrate Age Gradient chronosequence. Boxplots show the distribution of individual replicate samples and points indicate the predicted mean ( $\pm$  95% CI) species richness or diversity for each treatment of the unfertilized control (C). Significant direct and interactive effects of N and P additions are indicated to the right of each panel, where +  $P < 0.1$ , \*  $P < 0.05$ , \*\*  $P < 0.01$  and \*\*\*  $P < 0.001$ .

**Table 1.** Results of permutational MANOVAs examining the effect of fertilizer addition treatments on the composition of (a) all fungi or (b) putative ErMF associated with *V. calycinum* at the 300-yr and 4.1-myrr sites in the Long Substrate Age Gradient chronosequence.

(a) All fungi				
Site	Treatment	F	$r^2$	P
Young (300 yr)	Full model	1.876	0.167	<0.001
	N	2.284	0.068	0.003
	P	2.717	0.081	0.001
	N:P	0.651	0.019	0.913
Old (4.1 myr)	Full model	1.767	0.159	<0.001
	N	1.588	0.047	0.020
	P	2.647	0.080	<0.001
	N:P	1.066	0.032	0.345
(b) Putative ErMF				
Site	Treatment	F	$r^2$	P
Young (300 yr)	Full model	1.787	0.160	0.007
	N	1.488	0.044	0.130
	P	3.283	0.099	<0.001
	N:P	0.549	0.016	0.891
Old (4.1 myr)	Full model	2.324	0.199	<0.001
	N	1.782	0.051	0.028
	P	3.592	0.103	<0.001
	N:P	1.600	0.046	0.060

bias against some of these taxa. We did recover some sequences of the common ErMF species *Oidiodendron maius*, though at fairly low prevalence, and primarily in the oldest site. However, *O. maius* is also commonly found as a free-living soil saprotroph (Rice and Currah 2006) and our aggressive surface sterilization of roots may have biased our sampling against taxa that thrive outside of roots under some conditions.

In summary, we have shown that the diversity of fungi associated with the roots of ericaceous plants increases throughout long-term soil development largely due to increasing richness and evenness of putative ErMF taxa. Our results contrast with

recent reports of a hump-shaped diversity pattern with increasing soil age for arbuscular mycorrhizal fungi (Krüger et al. 2015), suggesting that functional differences among types of mycorrhizal fungi may determine the response of root-associated fungi to long-term soil development. We have also shown that P availability can affect the species composition and community structure of ericoid fungi, particularly in older P-limited ecosystems. Our results suggest that further study of ErMF from P-limited systems or across long-term soil chronosequences may prove valuable for identifying the extent of taxonomic and functional diversity for this unique group of mycorrhizal fungi.

## ACKNOWLEDGMENTS

We thank Michael Shintaku at the College of Agriculture, Forestry and Natural Resource Management and Anne Veillet at the Evolutionary Genomics Core Facility, both at the University of Hawaii, Hilo, for access to laboratory space and equipment. We also thank Matt Knope for feedback on earlier drafts of this paper. Site access was granted by the Hawaii Division of Land and Natural Resources, the Hawaii Experimental Tropical Forest and Parker Ranch.

## SUPPLEMENTARY DATA

Supplementary data are available at [FEMSEC](https://academic.oup.com/femsec) online.

## FUNDING

This work was supported by the Department of Biology at Stanford University and student research grants from the American Society of Naturalists and the Mycological Society of America.

**Conflicts of interest.** None declared.

## REFERENCES

- Abarenkov K, Zirk A, Piirmann T et al. UNITE general FASTA release for Fungi. 2020. DOI: 10.15156/BIO/786369.
- Adamczyk B, Ahvenainen A, Sietiö O-M et al. The contribution of ericoid plants to soil nitrogen chemistry and organic matter decomposition in boreal forest soil. *Soil Biol Biochem* 2016;**103**:394–404.
- Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 2001;**26**:32–46.
- Bruzzone MC, Fehrer J, Fontenla SB et al. First record of *Rhizoscyphus ericae* in Southern Hemisphere's Ericaceae. *Mycorrhiza* 2017;**27**:147–63.
- Cairney JWG, Meharg AA. Ericoid mycorrhiza: a partnership that exploits harsh edaphic conditions. *Eur J Soil Sci* 2003;**54**:735–40.
- Cairney JWG, Sawyer NA, Sharples JM et al. Intraspecific variation in nitrogen source utilisation by isolates of the ericoid mycorrhizal fungus *Hymenoscyphus ericae* (Read) Korf and Kernan. *Soil Biol Biochem* 2000;**32**:1319–22.
- Callahan BJ, McMurdie PJ, Rosen MJ et al. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016;**13**:581–3.
- Chadwick OA, Derry LA, Vitousek PM et al. Changing sources of nutrients during four million years of ecosystem development. *Nature* 1999;**397**:491–7.
- Chao A, Gotelli NJ, Hsieh TC et al. Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. *Ecol Monogr* 2014;**84**:45–67.
- Clemmensen KE, Bahr A, Ovaskainen O et al. Roots and associated fungi drive long-term carbon sequestration in boreal forest. *Science* 2013;**339**:1615–8.
- Clemmensen KE, Finlay RD, Dahlberg A et al. Carbon sequestration is related to mycorrhizal fungal community shifts during long-term succession in boreal forests. *New Phytol* 2015;**205**:1525–36.
- Cline LC, Zak DR. Dispersal limitation structures fungal community assembly in a long-term glacial chronosequence. *Environ Microbiol* 2014;**16**:1538–48.
- Crews TE, Kitayama K, Fownes JH et al. Changes in soil phosphorus fractions and ecosystem dynamics across a long chronosequence in Hawaii. *Ecology* 1995;**76**:1407–24.
- DeForest JL, Scott LG. Available organic soil phosphorus has an important influence on microbial community composition. *Soil Sci Soc Am J* 2010;**74**:2059.
- Dickie IA, Martínez-García LB, Koele N et al. Mycorrhizas and mycorrhizal fungal communities throughout ecosystem development. *Plant Soil* 2013;**367**:11–39.
- Dufrène M, Legendre P. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol Monogr* 1997;**67**:345–66.
- Fehrer J, Réblová M, Bambasová V et al. The root-symbiotic *Rhizoscyphus ericae* aggregate and *Hyaloscypha* (Leotiomyces) are congeneric: phylogenetic and experimental evidence. *Stud Mycol* 2019;**92**:195–225.
- Fukami T, Wardle DA. Long-term ecological dynamics: reciprocal insights from natural and anthropogenic gradients. *Proc Biol Sci* 2005;**272**:2105–15.
- Gillespie RG. Island time and the interplay between ecology and evolution in species diversification. *Evol Appl* 2016;**9**:53–73.
- Goh KM, Rafter TA, Stout JD et al. The accumulation of soil organic matter and its carbon isotope content in a chronosequence of soils developed on aeolian sand in New Zealand. *J Soil Sci* 1976;**27**:89–100.
- Grelet G-A, Johnson D, Paterson E et al. Reciprocal carbon and nitrogen transfer between an ericaceous dwarf shrub and fungi isolated from *Piceirhiza bicolorata* ectomycorrhizas. *New Phytol* 2009a;**182**:359–66.
- Grelet G-A, Meharg AA, Alexander IJ. Carbon availability affects nitrogen source utilisation by *Hymenoscyphus ericae*. *Mycol Res* 2005;**109**:469–77.
- Grelet G-A, Meharg AA, Duff EI et al. Small genetic differences between ericoid mycorrhizal fungi affect nitrogen uptake by *Vaccinium*. *New Phytol* 2009b;**181**:708–18.
- Hambleton S, Sigler L. *Meliniomyces*, a new anamorph genus for root-associated fungi with phylogenetic affinities to *Rhizoscyphus ericae* (= *Hymenoscyphus ericae*), Leotiomyces. *Stud Mycol* 2005;**53**:1–27.
- Harrington RA, Fownes JH, Vitousek PM. Production and resource use efficiencies in N- and P-limited tropical forests: a comparison of responses to long-term fertilization. *Ecosystems* 2001;**4**:646–57.
- Hazard C, Gosling P, Mitchell DT et al. Diversity of fungi associated with hair roots of ericaceous plants is affected by land use. *FEMS Microbiol Ecol* 2014;**87**:586–600.
- Herbert Da, Fownes JH. Phosphorus limitation of forest leaf area and net primary production on a highly weathered soil. *Biogeochemistry* 1995;**29**:223–35.
- Hobbie SE, Vitousek PM. Nutrient limitation of decomposition in Hawaiian forests. *Ecology* 2000;**81**:1867–77.
- Hsieh TC, Ma KH, Chao A. iNEXT: interpolation and extrapolation for species diversity. R package version 2.0.8. R-project 2016:1–18. DOI: 10.1111/2041-210X.12613.
- Hutton BJ, Dixon KW, Sivasithamparam K et al. Effect of habitat disturbance on inoculum potential of ericoid endophytes of Western Australian heaths (Epacridaceae). *New Phytol* 1997;**135**:739–44.
- Ishida TA, Nordin A. No evidence that nitrogen enrichment affect fungal communities of *Vaccinium* roots in two contrasting boreal forest types. *Soil Biol Biochem* 2010;**42**:234–43.
- Johnson NC, Wilson GWT, Bowker MA et al. Resource limitation is a driver of local adaptation in mycorrhizal symbioses. *Proc Natl Acad Sci* 2010;**107**:2093–8.
- Kiers ET, Duhamel M, Beesetty Y et al. Reciprocal rewards stabilize cooperation in the mycorrhizal symbiosis. *Science* 2011;**333**:880–2.
- Kitayama K, Mueller-Dombois D. Vegetation changes along gradients of long-term soil development in the Hawaiian montane rainforest zone. *Vegetatio* 1995;**120**:1–20.
- Kohout P. Biogeography of ericoid mycorrhiza. In: Tedersoo L (ed.). *Biogeography of Mycorrhizal Symbiosis*. Springer, 2017, 179–93.
- Kolařík M, Vohník M. When the ribosomal DNA does not tell the truth: the case of the taxonomic position of *Kurtia argillacea*, an ericoid mycorrhizal fungus residing among Hymenochaetales. *Fungal Biol* 2018;**122**:1–18.
- Konvalinková T, Püschel D, Řezáčová V et al. Carbon flow from plant to arbuscular mycorrhizal fungi is reduced under phosphorus fertilization. *Plant Soil* 2017;**419**:319–33.
- Koske RE, Gemma JN, Englander L. Vesicular-arbuscular mycorrhizae in Hawaiian Ericales. *Am J Bot* 1990;**77**:64–68.
- Koske RE, Gemma JN, Flynn T. Mycorrhizae in Hawaiian angiosperms: a survey with implications for the origin of the native flora. *Am J Bot* 1992;**79**:853–62.
- Krüger M, Teste FP, Laliberté E et al. The rise and fall of arbuscular mycorrhizal fungal diversity during ecosystem retrogression. *Mol Ecol* 2015;**24**:4912–30.

- Lambers H, Raven JA, Shaver GR et al. Plant nutrient-acquisition strategies change with soil age. *Trends Ecol Evol* 2008;**23**: 95–103.
- Leake JR, Read DJ. Experiments with ericoid mycorrhiza. *Methods Microbiol* 1991;**23**:435–57.
- Leopold DR, Fukami T. Greater local diversity under older species pools may arise from enhanced competitive equivalence. *Ecol Lett* 2020;**24**:310–18.
- Leopold DR. Ericoid fungal diversity: challenges and opportunities for mycorrhizal research. *Fungal Ecol* 2016;**24**:114–23.
- Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetjournal* 2011;**17**:10.
- Martínez-García LB, Richardson SJ, Tylianakis JM et al. Host identity is a dominant driver of mycorrhizal fungal community composition during ecosystem development. *New Phytol* 2015;**205**:1565–76.
- McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. Watson M (ed.). *PLoS One* 2013;**8**:e61217.
- McMurdie PJ, Holmes S. Waste not, want not: why rarefying microbiome data is inadmissible. McHardy AC (ed.). *PLoS Comput Biol* 2014;**10**:e1003531.
- Mitchell DT, Gibson BR. Ericoid mycorrhizal association: ability to adapt to a broad range of habitats. *Mycologist* 2006;**20**:2–9.
- Myers MD, Leake JR. Phosphodiesterases as mycorrhizal P sources II. Ericoid mycorrhiza and the utilization of nuclei as a phosphorus and nitrogen source by *Vaccinium macrocarpon*. *New Phytol* 1996;**132**:445–51.
- Nasholm T, Ekblad A, Nordin A et al. Boreal forest plants take up organic nitrogen. *Nature* 1998;**392**:914–6.
- Newsham KK. A meta-analysis of plant responses to dark septate root endophytes. *New Phytol* 2011;**190**:783–93.
- Oksanen J, Blanchet FG, Friendly M et al. vegan: Community Ecology Package. R package version 2.5-7. 2017:1–292. URL:<https://CRAN.R-project.org/package=vegan> (January 2020, date last accessed).
- Olsson PA, Rahm J, Aliasghar zad N. Carbon dynamics in mycorrhizal symbioses is linked to carbon costs and phosphorus benefits. *FEMS Microbiol Ecol* 2010;**72**:125–31.
- Ostertag R, Verville JH. Fertilization with nitrogen and phosphorus increases abundance of non-native species in Hawaiian montane forests. *Plant Ecol* 2002;**62**:77–90.
- Pearson V, Read DJ. Biology of mycorrhiza in the Ericaceae I. The isolation of the endophyte and synthesis of mycorrhizas in aseptic culture. *New Phytol* 1973a;**72**: 371–9.
- Pearson V, Read DJ. Biology of mycorrhiza in the Ericaceae II. The transport of carbon and phosphorus by the endophyte and the mycorrhiza. *New Phytol* 1973b;**72**:1325–31.
- Peltzer DA, Wardle DA, Allison VJ et al. Understanding ecosystem retrogression. *Ecol Monogr* 2010;**80**:509–29.
- Perotto S, Daghino S, Martino E. Ericoid mycorrhizal fungi and their genomes: another side to the mycorrhizal symbiosis? *New Phytol* 2018;**220**:1141–7.
- Perotto S, Martino E, Abba S et al. Genetic diversity and functional aspects of ericoid mycorrhizal fungi. In: Hock B (ed.). *Fungal Associations*. 2nd edn Springer Berlin Heidelberg, 2012, 255–85.
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/> (January 2020, date last accessed) 2020.
- Read DJ. Mycorrhizas in ecosystems. *Experientia* 1991;**47**:376–91.
- Read DJ. The biology of mycorrhiza in the Ericales. *Can J Bot* 1983;**61**:985–1004.
- Reed SC, Vitousek PM, Cleveland CC. Are patterns in nutrient limitation belowground consistent with those aboveground: results from a 4 million year chronosequence. *Biogeochemistry* 2011;**106**:323–36.
- Rice AV, Currah RS. *Oidiodendron maius*: saprobe in *Sphagnum* peat, mutualist in ericaceous roots? In: Schulz BJE, Boyle CJC, Sieber TN (eds.). *Microbial Root Endophytes*. Berlin, Heidelberg: Springer, 2006, 227–46.
- Richardson SJ, Peltzer DA, Allen RB et al. Rapid development of phosphorus limitation in temperate rainforest along the Franz Josef soil chronosequence. *Oecologia* 2004;**139**:267–76.
- Roberts DW. labdsv: Ordination and Multivariate Analysis for Ecology. R package version 2.0-1, 2019. URL: <https://CRAN.R-project.org/package=labdsv> (January 2020, date last accessed).
- Selosse M-A, Setaro S, Glatard F et al. Sebacinaleae are common mycorrhizal associates of Ericaceae. *New Phytol* 2007;**174**:864–78.
- Smith DP, Peay KG. Sequence depth, not PCR replication, improves ecological inference from next generation DNA sequencing. *PLoS One* 2014;**9**:e90234.
- Smith S, Read D. *Mycorrhizal Symbiosis*. 3rd edn Academic Press, 2006.
- Tamm CO. *Nitrogen in Terrestrial Ecosystems*. Berlin, Heidelberg: Springer Berlin Heidelberg, 1991.
- Tisserant E, Malbreil M, Kuo A et al. Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. *Proc Natl Acad Sci* 2013;**110**:20117–22.
- Torn MS, Trumbore SE, Chadwick OA et al. Mineral control of soil organic carbon storage and turnover. *Nature* 1997;**389**:170–3.
- Treseder K, Allen M. Direct nitrogen and phosphorus limitation of arbuscular mycorrhizal fungi: a model and field test. *New Phytol* 2002;**155**:507–15.
- Turner BL. Resource partitioning for soil phosphorus: a hypothesis. *J Ecol* 2008;**96**:698–702.
- Van Geel M, Jacquemyn H, Peeters G et al. Diversity and community structure of ericoid mycorrhizal fungi in European bogs and heathlands across a gradient of nitrogen deposition. *New Phytol* 2020, DOI: 10.1111/nph.16789.
- Villarreal-Ruiz L, Anderson IC, Alexander IJ. Interaction between an isolate from the *Hymenoscyphus ericae* aggregate and roots of *Pinus* and *Vaccinium*. *New Phytol* 2004;**164**:183–92.
- Villarreal-Ruiz L, Neri-Luna C, Anderson IC et al. In vitro interactions between ectomycorrhizal fungi and ericaceous plants. *Symbiosis* 2012;**56**:67–75.
- Vitousek P. *Nutrient Cycling and Limitations: Hawaii as a Model System*. Princeton University Press, 2004.
- Vitousek PM, Farrington H. Nutrient limitation and soil development: experimental test of a biogeochemical theory. *Biogeochemistry* 1997;**37**:63–75.
- Vitousek PM, Howarth RW. Nitrogen limitation on land and in the sea: how can it occur? *Biogeochemistry* 1991;**13**:87–115.
- Vitousek PM, Walker LR, Whiteaker LD et al. Nutrient limitations to plant growth during primary succession in Hawaii Volcanoes National Park. *Biogeochemistry* 1993;**23**:197–215.
- Vohník M, Mrnka L, Lukešová T et al. The cultivable endophytic community of Norway spruce ectomycorrhizas from microhabitats lacking ericaceous hosts is dominated by ericoid mycorrhizal *Meliniomyces variabilis*. *Fungal Ecol* 2013;**6**:281–92.
- Vohník M, Pánek M, Fehrer J et al. Experimental evidence of ericoid mycorrhizal potential within Serendipitaceae (Sebaciniales). *Mycorrhiza* 2016;**26**:831–46.
- Vohník M, Sadowsky JJ, Kohout P et al. Novel root-fungus symbiosis in Ericaceae: sheathed ericoid mycorrhiza formed

- by a hitherto undescribed basidiomycete with affinities to Trechisporales. *PLoS One* 2012;7:e39524.
- Vrålstad T, Fossheim T, Schumacher T. *Piceirhiza bicolorata* - the ectomycorrhizal expression of the *Hymenoscyphus ericae* aggregate? *New Phytol* 2000;145:549–63.
- Vrålstad T, Myhre E, Schumacher T. Molecular diversity and phylogenetic affinities of symbiotic root-associated ascomycetes of the Helotiales in burnt and metal polluted habitats. *New Phytol* 2002;155:131–48.
- Vrålstad T. Are ericoid and ectomycorrhizal fungi part of a common guild? *New Phytol* 2004;164:7–10.
- Walker LR, Wardle DA, Bardgett RD et al. The use of chronosequences in studies of ecological succession and soil development. *J Ecol* 2010;98:725–36.
- Walker TW, Syers JK. The fate of phosphorus during pedogenesis. *Geoderma* 1976;15:1–19.
- Wang Q, Garrity GM, Tiedje JM et al. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 2007;73:5261–7.
- Wardle DA. Ecosystem properties and forest decline in contrasting long-term chronosequences. *Science* 2004;305:509–13.
- Whittaker S, Cairney J. Influence of amino acids on biomass production by ericoid mycorrhizal endophytes from *Woollisia pungens* (Epacridaceae). *Mycol Res* 2001;105:105–11.
- Wickham H, Averick M, Bryan J et al. Welcome to the Tidyverse. *J Open Source Softw* 2019;4:1686.
- Wright ES. Using DECIPHER v2.0 to analyze big biological sequence data in R. *R J* 2016;8:352.
- Wurzburger N, Higgins BP, Hendrick RL. Ericoid mycorrhizal root fungi and their multicopper oxidases from a temperate forest shrub. *Ecol Evol* 2012;2:65–79.
- Xiao G, Berch SM. Organic nitrogen use by salal ericoid mycorrhizal fungi from northern Vancouver Island and impacts on growth in vitro of *Gaultheria shallon*. *Mycorrhiza* 1999;9:145–9.