

International Journal of Game Theory

A Publication of the Institute for Advanced Studies, Vienna

Founded by Oskar Morgenstern

Editorial Board

- Robert J. Aumann, Hebrew University of Jerusalem
Claude Berge, University of Paris
Louis J. Billera, Cornell University, Ithaca
Karl H. Borch, Norwegian School of Economics and Business Administration, Bergen
John C. Harsanyi, University of California, Berkeley
Michael Maschler, Hebrew University of Jerusalem
Guillermo Owen, Instituto SER, Bogota
Bezalel Peleg, Hebrew University of Jerusalem
Anatol Rapoport, University of Toronto
Robert Reichardt, University of Vienna
Reinhard Selten, University of Bielefeld
Lloyd S. Shapley, RAND Corporation, Santa Monica
Martin Shubik, Yale University, New Haven
Patrick Suppes, Stanford University
Mitsuo Suzuki, Tokyo Institute of Technology
Robert M. Thrall, Rice University, Houston
N.N. Vorob'ev, Academy of Sciences, Leningrad

Scientific Editor

William F. Lucas, Cornell University, Ithaca

Managing Editor

Gerhard Schwödiauer, University of Bielefeld



Physica-Verlag · Vienna

Volume 8

1979

Issue 3

CONTENTS

<i>Roth, A.E.</i> : An Impossibility Result Concerning n -Person Bargaining Games	129
<i>Shenoy, P.P.</i> : On Coalition Formation: A Game-Theoretical Approach	133
<i>Muto, S.</i> : Generalized k -Quota Solutions for (N, k) Games	165
<i>Frick, H.</i> : On an Application of Game Theory to a Problem of Testing Statistical Hypothesis	175

SCIENTIFIC EDITOR:

Prof. Dr. W.F. Lucas
Cornell University
334 Upson Hall
Ithaca, N.Y. 14853/USA

MANAGING EDITOR:

Dr. Gerhard Schwödauer
University of Bielefeld
Faculty of Economics
POB 8640
D-4800 Bielefeld 1

The *International Journal of Game Theory* appears quarterly. It publishes original contributions to game theory with substantial mathematical content. Occasionally, other papers may also be published if they are of considerable interest to game theorists.

Besides original contributions, the *International Journal of Game Theory* offers a "Listing Service" which announces research memoranda, discussion papers etc. with the names and addresses of the authors. Authors of such papers appearing elsewhere, are asked to send a copy or offprint to the Scientific Editor.

The annual subscription rate is DM 130.- resp. S. 925.- + postage.

Orders can be placed directly with the publisher. All orders must be accompanied by payment. Orders from German booksellers may be placed with Physica-Verlag, Würzburg.

The *International Journal of Game Theory* or parts thereof, may not be translated or reproduced in any form without written permission of the publisher.

Composed and printed by: „Journalfranz“ Arnulf Liebing GmbH + Co., Würzburg



PHYSICA-VERLAG Rudolf Liebing KG • Seilerstätte 18, A-1010 Wien

1979

An Impossibility Result Concerning n -Person Bargaining Games¹⁾

By *A.E. Roth*, Urbana²⁾

Abstract: In this note we show that a solution proposed by *Raiffa* for two-person bargaining games, which has recently been axiomatized by *Kalai/Smorodinsky*, does not generalize in a straightforward manner to general n -person bargaining games. Specifically, the solution is not Pareto optimal on the class of all n -person bargaining games, and no solution which is can possess the other properties which characterize *Raiffa's* solution in the two-person case.

An n -person bargaining game consists of a set of players $N = \{1, \dots, n\}$, a vector of utilities $d = (d_1, \dots, d_n)$, and a compact convex subset $S \subset R^n$ of utility vectors which contains the vector d and at least one element x such that $x > d$. The interpretation is that the set S is the set of feasible expected utility payoffs to the players, any one of which can be achieved by unanimous agreement. In the event that no unanimous agreement is reached, the disagreement vector d is the result.³⁾

A solution is a function defined on the class B of such bargaining games, which selects a feasible outcome for every game: i.e., $f: B \rightarrow R^n$ such that for any game (S, d) in B , $f(S, d) = x$ is an element of S . *Nash* [1950] was the first to study solutions to the bargaining game, which he interpreted as modelling the anticipated outcome of the game when it is played by rational players. *Raiffa* [1953] also studied solutions, which he interpreted as indicating the manner in which games might be arbitrated by an impartial arbiter.

Nash approached the problem axiomatically, and specified a list of properties which collectively characterize a unique solution. Although *Nash* explicitly studied only the case of two-person bargaining his treatment generalizes without any difficulty to the general n -person case.⁴⁾ *Nash* required that a solution possess the following properties.⁵⁾

¹⁾ This work was supported by National Science Foundation Grant SOC 78-09928 to the University of Illinois. It has also benefitted from discussion and correspondence with H. Imai, E. Kalai, and M. Maschler.

²⁾ Prof. *A.E. Roth*, Department of Business Administration, University of Illinois, Urbana, Ill. 61801, USA.

³⁾ That is, if the players unanimously agree on some x in S , then each player i receives the utility x_i . If no agreement is reached, player i receives the utility d_i .

⁴⁾ The two-person case is the one which has been studied the most, both because of its simplicity, as well as because most two-person cooperative games can be thought of as bargaining games, whereas bargaining games are only a subclass of games with more than two players.

⁵⁾ These properties are stated here in a slightly different form than in *Nash's* original paper.

Property 1. Pareto optimality: If $f(S, d) = x$, then there exists no y in S , distinct from x , such that $y \geq x$.

Property 2. Symmetry: If (S, d) is a symmetric game,⁶) then $f_i(S, d) = f_j(S, d)$ for all i, j in N .

Property 3. Independence of equivalent utility representations: Let $T: R^n \rightarrow R^n$ be an order-preserving linear transformation of the players' utility functions (i.e., if $y = T(x)$, then $y_i = a_i x_i + b_i$, where $a_i > 0$). Then $f(T(S), T(d)) = T(f(S, d))$.

Property 4. Independence of irrelevant alternatives: If T contains S , and $f(T, d) \in S$, then $f(S, d) = f(T, d)$.

These properties have been extensively discussed in the literature [cf. *Luce/Raiffa; Harsanyi; Kalai*, 1977a; *Roth*, 1977b, 1978, 1979b], and so we will not review that discussion here.

Nash proved that there is a unique solution which possesses properties 1–4. It is the solution $f = F$ defined by $F(S, d) = x$ such that $x > d$ and $\Pi(x_i - d_i) \geq \Pi(y_i - d_i)$ for all y in S such that $y > d$.

That is, *Nash's* solution selects the strongly individually rational outcome which maximizes the geometric average of the gains available to the players. Formally, this result can be stated as follows.

Theorem 1: F is the unique solution which is Pareto optimal, symmetric, independent of equivalent utility representations, and independent of irrelevant alternatives.

It has recently been shown [*Roth*, 1977a] that the property of strong individual rationality (i.e., $f(S, d) > d$) can replace Pareto optimality in the characterization of *Nash's* solution.

Since *Nash's* solution is independent of irrelevant alternatives, it is not sensitive to the range of outcomes contained in the feasible set, as reflected, for instance, by the ideal point \bar{x} defined by $\bar{x}_i = \max\{x_i \mid x \in S \text{ and } x \geq d\}$ for $i \in N$. The point \bar{x} can be thought of as reflecting the potential aspirations of the players, and *Raiffa* [1953] proposed a solution for two-player games which is sensitive to changes in \bar{x} .⁷) Specifically, *Raiffa* proposed the solution G for two-player games such that $G(S, d) = x$ is the Pareto optimal point at which $(x_1 - d_1)/(x_2 - d_2) = (\bar{x}_1 - d_1)/(\bar{x}_2 - d_2)$. That is, the solution G selects the maximal point on the line joining d to \bar{x} , yielding each player the largest reward consistent with the constraint that the players' actual gains should be in proportion to their potential gains, as measured by the ideal point \bar{x} .

Kalai/Smorodinsky [1975] give an elegant axiomatization of the solution G , and show that it is the unique solution defined on the class of two-person games which

⁶) A game (S, d) is symmetric if $d_i = d_j$ for all i, j in N , and if for every element x in S , every permutation $y = \pi x$ is also contained in S .

⁷) The same solution has been proposed by *Crott* [1971], on the basis of some experimental results, and it has also been studied by *Butrim* [1976], who attributes it to the Russian game-theorist *Germeier*.

possesses properties 1–3 as well as the following property.⁸⁾

Property 5: Restricted monotonicity: Let (T, d) and (S, d) be games which share a common ideal point, and such that T contains S . Then $f(T, d) \geq f(S, d)$.

That is, *Kalai/Smorodinsky* proved the following.

Theorem 2: The solution G is the unique solution defined on the class of two-person games which possesses properties 1, 2, 3, and 5.

In view of the fact that *Nash's* solution is so easily generalized to bargaining games with more than two players, we might expect that this would also be the case for the solution G . Instead, it turns out that for games with more than two players, the solution G no longer possesses all the properties named in the theorem, and there exists no solution which does. We will see that the solution G fails to select a Pareto optimal outcome for all n person games with $n \geq 3$, although it continues to obey the other properties named in Theorem 2. In fact, for games with more than two players, there exists no solution defined on the class B which possesses properties 1, 2, and 5. Formally, we will prove the following.

Theorem 3: For bargaining games with three or more players, no solution exists which possesses the properties of Pareto optimality, symmetry, and restricted monotonicity.

Proof: We will assume that f is a solution which possesses the properties named in the theorem, and show that this leads to a contradiction. Consider the n -person game ($n \geq 3$) whose disagreement point is equal to the origin (which we will denote by \bar{O}), and whose feasible set S is equal to the convex hull of \bar{O} and the points p and q such that $p_1 = 0$ and $p_i = 1$ for $i \neq 1$, and $q_2 = 0$ and $q_j = 1$ for $j \neq 2$. Then the set of Pareto optimal points in S is the line segment joining p to q , so $f(S, \bar{O}) = x$ is a convex combination of p and q . In particular, $x_3 = 1$.

Now consider the game (T, \bar{O}) , where $T = \{x \geq \bar{O} \mid \sum x_i \leq (n-1) \text{ and } x_i \leq 1 \text{ for } i = 1, \dots, n\}$. Then (T, \bar{O}) is a symmetric game, so $f_1(T, \bar{O}) = f_2(T, \bar{O}) = \dots = f_n(T, \bar{O})$, and the Pareto optimality of f implies that

$$f(T, \bar{O}) = z = \left(\frac{(n-1)}{n}, \frac{(n-1)}{n}, \dots, \frac{(n-1)}{n} \right).$$

But the point $(1, 1, \dots, 1)$ is the ideal point both of the game (S, \bar{O}) and of the game (T, \bar{O}) , so the fact that T contains S and the restricted monotonicity of f imply that $z \geq x$. Since we have shown that $z_3 = (n-1)/n$ and $x_3 = 1$, this gives the contradiction needed to complete the proof.

⁸⁾ Actually, *Kalai/Smorodinsky* consider a stronger property which they call individual monotonicity, and which implies the property of restricted monotonicity stated here. However their treatment makes it clear that only this weaker condition is necessary to characterize the solution G for two-person games.

⁹⁾ Note that the solution G is not even weakly Pareto optimal, as it would be if we confined our attention to games with disposable utility. (For a discussion of some matters related to this point, see *Kalai* [1977b], *Myerson* [1977], and *Roth* [1979a]).

Note that the solution G does not fail to be well-defined for $n \geq 3$, since for any game (S, d) there is still a unique maximal point on the line joining d to the ideal point \bar{x} . Furthermore, it is straightforward to confirm that the solution G possesses properties 2, 3, and 5 for any n . Theorem 3 therefore implies that G must not possess property 1, and in fact, in the game (S, \bar{O}) considered in the proof of the theorem, $G(S, \bar{O}) = \bar{O}$, which is not Pareto optimal.⁹

These results suggest that there may be some fundamental differences between the general case and the two-person case, and that other solution concepts will need to be explored in any attempt to incorporate players' aspirations in a consistent way into multi-player games.

References

- Butrim, B.I.*: A Modified Solution of the Bargaining Problem. (Russian), Zhurnal Vychislitel' noi Matematiki Matematicheskoi Fiziki 16, 1976, 340–350.
- Crott, H.W.*: Experimentelle Untersuchung zum Verhandlungsverhalten in Kooperativen Spielen. Zeitschrift für Sozialpsychologie 2, 1971, 61–74.
- Harsanyi, J.C.*: Rational Behavior and Bargaining Equilibrium in Games and Social Situations. Cambridge 1977.
- Kalai, E.*: Nonsymmetric Nash Solutions and Replications of 2-Person Bargaining. International Journal of Game Theory, 1977a, 129–133.
- : Proportional Solutions to Bargaining Situations: Interpersonal Utility Comparison. Econometrica 45, 1977b, 1623–1630.
- Kalai, E., and M. Smorodinsky*: Other Solutions to Nash's Bargaining Problem. Econometrica 43, 1975, 513–518.
- Luce, R.D., and H. Raiffa*: Games and Decisions: Introduction and Critical Survey. New York 1957.
- Myerson, R.B.*: Two-Person Bargaining Problems and Comparable Utility. Econometrica 45, 1977, 1631–1637.
- Nash, J.F.*: The Bargaining Problem. Econometrica 28, 1950, 155–162.
- Raiffa, H.*: Arbitration Schemes for Generalized Two-Person Games. Contributions to the Theory of Games II. Ed. by H.W. Kuhn and A.W. Tucker. Annals of Mathematics Studies, No. 28, Princeton 1953, 361–387.
- Roth, A.E.*: Individual Rationality and Nash's Solution to the Bargaining Problem. Mathematics of Operations Research 2, 1977a, 64–65.
- : Independence of Irrelevant Alternatives and Solutions to Nash's Bargaining Problem. Journal of Economic Theory 16, 1977b, 247–251.
- : The Nash Solution and the Utility of Bargaining. Econometrica 46, 1978, 587–594, 983.
- : Proportional Solutions to the Bargaining Problems. Econometrica 47, 1979a, 775–778.
- : Axiomatic Models of Bargaining. Lecture Notes in Economics and Mathematical Systems, # 170, Berlin–Heidelberg–New York, forthcoming, 1979b.

Received September, 1978
(revised version July, 1979)