

Math Camp

Justin Grimmer

Associate Professor
Department of Political Science
Stanford University

September 6th, 2016

Lab this afternoon!

130-300pm

Convergence

Big idea today is **convergence**

Convergence

Big idea today is **convergence**

- **Sequence** \rightarrow converge on some number

Convergence

Big idea today is **convergence**

- **Sequence** → converge on some number
- **Function** → **limit** (use to calculate derivatives)

Convergence

Big idea today is **convergence**

- **Sequence** → converge on some number
- **Function** → **limit** (use to calculate derivatives)
- **Continuity** → a function doesn't jump (converge on itself)

Convergence

Big idea today is **convergence**

- **Sequence** → converge on some number
- **Function** → **limit** (use to calculate derivatives)
- **Continuity** → a function doesn't jump (converge on itself)
- **Derivatives** → limits that measure a function's properties

Sequence: Definition + Examples

Definition


A *sequence* is a function whose domain is the set of positive integers

We'll write a sequence as,

$$\{a_n\}_{n=1}^{\infty} = (a_1, a_2, \dots, a_N, \dots)$$


Sequence: Definition + Examples

$$\left\{ \frac{1}{n} \right\} = (1, 1/2, 1/3, 1/4, \dots, 1/N, \dots)$$




Sequence: Definition + Examples

$$\left\{ \frac{1}{n^2} \right\} = (1, 1/4, 1/9, 1/16, \dots, 1/N^2, \dots)$$




Sequence: Definition + Examples

$$\left\{ \frac{1 + (-1)^n}{2} \right\} = (0, 1, 0, 1, \dots, 0, 1, 0, 1, \dots)$$



Sequence: Definition + Examples


$$\{\theta\}_{n=1}^{\infty} = (\theta_1, \theta_2, \dots, \theta_n, \dots)$$
$$\theta_n = f(\text{n responses (vote choice)})$$



Sequence: Convergence

Consider the sequence


$$\left\{ \frac{(-1)^n}{n} \right\} = \left(-1, \frac{1}{2}, \frac{-1}{3}, \frac{1}{4}, \frac{-1}{5}, \frac{1}{6}, \frac{-1}{7}, \frac{1}{8}, \dots \right)$$



Sequence: Convergence

Consider the sequence


$$\left\{ \frac{(-1)^n}{n} \right\} = \left(-1, \frac{1}{2}, \frac{-1}{3}, \frac{1}{4}, \frac{-1}{5}, \frac{1}{6}, \frac{-1}{7}, \frac{1}{8}, \dots \right)$$



Sequence: Convergence

Consider the sequence


$$\left\{ \frac{(-1)^n}{n} \right\} = \left(-1, \frac{1}{2}, \frac{-1}{3}, \frac{1}{4}, \frac{-1}{5}, \frac{1}{6}, \frac{-1}{7}, \frac{1}{8}, \dots \right)$$



Sequence: Convergence

Consider the sequence


$$\left\{ \frac{(-1)^n}{n} \right\} = \left(-1, \frac{1}{2}, \frac{-1}{3}, \frac{1}{4}, \frac{-1}{5}, \frac{1}{6}, \frac{-1}{7}, \frac{1}{8}, \dots \right)$$



Sequence: Convergence

Consider the sequence

$$\left\{ \frac{(-1)^n}{n} \right\} = \left(-1, \frac{1}{2}, \frac{-1}{3}, \frac{1}{4}, \frac{-1}{5}, \frac{1}{6}, \frac{-1}{7}, \frac{1}{8}, \dots \right)$$



Sequence: Convergence definition

Definition

A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a real number A if for each $\epsilon > 0$ there is a positive integer N such that for all $n \geq N$ we have $|a_n - A| < \epsilon$

Sequence: Convergence definition

Definition

A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a real number A if for each $\epsilon > 0$ there is a positive integer N such that for all $n \geq N$ we have $|a_n - A| < \epsilon$

- 1) If a sequence converges, it converges to **one** number. We call that **A**

Sequence: Convergence definition

Definition

A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a real number A if for each $\epsilon > 0$ there is a positive integer N such that for all $n \geq N$ we have $|a_n - A| < \epsilon$

- 1) If a sequence converges, it converges to **one** number. We call that **A**
- 2) $\epsilon > 0$ is some **arbitrary** real-valued number.

Sequence: Convergence definition

Definition

A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a real number A if for each $\epsilon > 0$ there is a positive integer N such that for all $n \geq N$ we have $|a_n - A| < \epsilon$

- 1) If a sequence converges, it converges to **one** number. We call that **A**
- 2) $\epsilon > 0$ is some **arbitrary** real-valued number. Think about this as our **error** tolerance. Notice $\epsilon > 0$.

Sequence: Convergence definition

Definition

A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a real number A if for each $\epsilon > 0$ there is a positive integer N such that for all $n \geq N$ we have $|a_n - A| < \epsilon$

- 1) If a sequence converges, it converges to **one** number. We call that **A**
- 2) $\epsilon > 0$ is some **arbitrary** real-valued number. Think about this as our **error** tolerance. Notice $\epsilon > 0$.
- 3) As we will see the N will depend upon ϵ


Sequence: Convergence definition

Definition


A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a real number A if for each $\epsilon > 0$ there is a positive integer N such that for all $n \geq N$ we have $|a_n - A| < \epsilon$

- 1) If a sequence converges, it converges to **one** number. We call that **A**
- 2) $\epsilon > 0$ is some **arbitrary** real-valued number. Think about this as our **error** tolerance. Notice $\epsilon > 0$.
- 3) As we will see the N will depend upon ϵ
- 4) Implies the sequence never gets further than ϵ away from A


Sequence: Convergence definition




Sequence: Convergence definition




Sequence: Convergence definition



Sequence: Convergence definition



Sequence: Convergence definition



Sequence: Proof of Convergence

Theorem

$\left\{\frac{1}{n}\right\}$ converges to 0

Proof.

We need to show that for ϵ there is some N_ϵ such that, for all $n \geq N_\epsilon$ $\left|\frac{1}{n} - 0\right| < \epsilon$. **Without loss of generality** (WLOG) select an ϵ . Then,

$$\begin{aligned}\left|\frac{1}{N_\epsilon} - 0\right| &< \epsilon \\ \frac{1}{N_\epsilon} &< \epsilon \\ \frac{1}{\epsilon} &< N_\epsilon\end{aligned}$$

For each epsilon, then, any $N_\epsilon > \frac{1}{\epsilon}$ will suffice. □

Sequence: Divergence + Bounded

Definition

*If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it bounded*

Sequence: Divergence + Bounded

Definition

*If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it bounded*

- An unbounded sequence

Sequence: Divergence + Bounded

Definition

*If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it bounded*

- An unbounded sequence

$$\{n\} = (1, 2, 3, 4, \dots, N, \dots)$$

Sequence: Divergence + Bounded

Definition

*If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it bounded*

- An unbounded sequence

$$\{n\} = (1, 2, 3, 4, \dots, N, \dots)$$

- A bounded sequence that doesn't converge

Sequence: Divergence + Bounded

Definition

If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it **bounded**

- An unbounded sequence

$$\{n\} = (1, 2, 3, 4, \dots, N, \dots)$$

- A bounded sequence that doesn't converge

$$\left\{ \frac{1 + (-1)^n}{2} \right\} = (0, 1, 0, 1, \dots, 0, 1, 0, 1, \dots)$$

Sequence: Divergence + Bounded

Definition

If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it **bounded**

- An unbounded sequence

$$\{n\} = (1, 2, 3, 4, \dots, N, \dots)$$

- A bounded sequence that doesn't converge

$$\left\{ \frac{1 + (-1)^n}{2} \right\} = (0, 1, 0, 1, \dots, 0, 1, 0, 1, \dots)$$

- **All convergent sequences are bounded**

Sequence: Divergence + Bounded

Definition

If a sequence, $\{a_n\}$ converges we'll call it **convergent**. If it doesn't we'll call it **divergent**. If there is some number M such that, for all n $|a_n| < M$, then we'll call it **bounded**

- An unbounded sequence

$$\{n\} = (1, 2, 3, 4, \dots, N, \dots)$$

- A bounded sequence that doesn't converge

$$\left\{ \frac{1 + (-1)^n}{2} \right\} = (0, 1, 0, 1, \dots, 0, 1, 0, 1, \dots)$$

- **All convergent sequences are bounded**
- If a sequence is **constant**, $\{C\}$ it converges to C . **proof?**

Algebra of Sequences

How do we add, multiply, and divide sequences?

Theorem

Suppose $\{a_n\}$ converges to A and $\{b_n\}$ converges to B . Then,

- $\{a_n + b_n\}$ converges to $A + B$
- $\{a_n b_n\}$ converges to $A \times B$.
- *Suppose $b_n \neq 0 \forall n$ and $B \neq 0$. Then $\left\{\frac{a_n}{b_n}\right\}$ converges to $\frac{A}{B}$.*

Working Together

- Consider the sequence $\left\{\frac{1}{n}\right\}$ —what does it converge to?
- Consider the sequence $\left\{\frac{1}{2n}\right\}$ what does it converge to?

Challenge Questions

- What does $\left\{3 + \frac{1}{n}\right\}$ converge to?
- What about $\left\{\left(3 + \frac{1}{n}\right)\left(100 + \frac{1}{n^4}\right)\right\}$?
- Finally, $\left\{\frac{300 + \frac{1}{n}}{100 + \frac{1}{n^4}}\right\}$?

Work smarter, not harder

Divide into teams, let's reconvene in about 10 minutes.

Sequences \rightsquigarrow Limits of Functions


Calculus/Real Analysis: study of functions on the **real line**.

Limit of a function: how does a function behave as it gets close to a particular point?

- Derivatives
- Asymptotics
- Game Theory


Limits of Functions

$$f(x) = \sin(x)$$




Limits of Functions

$$f(x) = \sin(x)$$




Limits of Functions

$$f(x) = \sin(x)$$




Limits of Functions

$$f(x) = \sin(x)$$




Limits of Functions

$$f(x) = \sin(x)$$




Limits of Functions

$$f(x) = \sin(x)$$



Limits of Functions

$$f(x) = \sin(x)$$



Precise Definition of Limits of Functions

Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. We say that f has a limit L at x_0 if, for each $\epsilon > 0$, there is a $\delta > 0$ such that $|x - x_0| < \delta$ implies that $|f(x) - L| < \epsilon$.

- Limits are about the behavior of functions at **points**. Here x_0 .
- As with sequences, we let ϵ define an **error rate**
- δ defines an area around x_0 where $f(x)$ is going to be within our error rate

Precise Definition of Limit: Example

Theorem

The function $f(x) = x + 1$ has a limit of 1 at $x_0 = 0$.

Proof.

WLOG choose $\epsilon > 0$. We want to show that there is δ_ϵ such that, $|x - x_0| < \delta_\epsilon$ implies $|f(x) - 1| < \epsilon$. In other words,

$$|x| < \delta_\epsilon \quad \text{implies} \quad |(x + 1) - 1| < \epsilon$$

$$|x| < \delta_\epsilon \quad \text{implies} \quad |x| < \epsilon$$

But if $\delta_\epsilon = \epsilon$ then this holds, we are done. □


Precise Definition of Limit: Example

A function can have a limit of L at x_0 even if $f(x_0) \neq L(!)$

Theorem

The function $f(x) = \frac{x^2-1}{x-1}$ has a limit of 2 at $x_0 = 1$.

$$f(x) = (x^2 - 1)/(x - 1)$$




Precise Definition of Limit: Example

A function can have a limit of L at x_0 even if $f(x_0) \neq L(!)$

Theorem

The function $f(x) = \frac{x^2-1}{x-1}$ has a limit of 2 at $x_0 = 1$.

$$f(x) = (x^2 - 1)/(x - 1)$$



Precise Definition of Limit: Example

Proof.

For all $x \neq 1$,

$$\begin{aligned}\frac{x^2 - 1}{x - 1} &= \frac{(x + 1)(x - 1)}{x - 1} \\ &= x + 1\end{aligned}$$

Choose $\epsilon > 0$ and set $x_0 = 1$. Then, we're looking for δ_ϵ such that

$$|x - 1| < \delta_\epsilon \quad \text{implies} \quad |(x + 1) - 2| < \epsilon$$


Again, if $\delta_\epsilon = \epsilon$, then this is satisfied.



Not all Functions have Limits!

Theorem

Consider $f : (0, 1) \rightarrow \mathbb{R}$, $f(x) = 1/x$. $f(x)$ does not have a limit at $x_0 = 0$



Proof.

Choose $\epsilon > 0$. We need to show that there **does not** exist δ such that

$$|x| < \delta \quad \text{implies} \quad \left| \frac{1}{x} - L \right| < \epsilon$$

But, there is a problem. Because

$$\begin{aligned} \frac{1}{x} - L &< \epsilon \\ \frac{1}{x} &< \epsilon + L \\ x &> \frac{1}{L + \epsilon} \end{aligned}$$

This implies that there **can't** be a δ , because x has to be bigger than $\frac{1}{L + \epsilon}$.

□

Intuitive Definition of Limit

Definition

If a function f tends to L at point x_0 we say it has a limit L at x_0 we commonly write,

$$\lim_{x \rightarrow x_0} f(x) = L$$

Definition

If a function f tends to L at point x_0 as we approach from the right, then we write

$$\lim_{x \rightarrow x_0^+} f(x) = L$$

and call this a **right hand limit**

If a function f tends to L at point x_0 as we approach from the left, then we write

$$\lim_{x \rightarrow x_0^-} f(x) = L$$

and call this a **left-hand limit**

Regression discontinuity designs

Left-hand, Right-hand, and Limits

Theorem

The $\lim_{x \rightarrow x_0} f(x)$ exists if and only if $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$

Left-hand, Right-hand, and Limits

Theorem

The $\lim_{x \rightarrow x_0} f(x)$ exists if and only if $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$

- Intuition that $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x) \Rightarrow \lim_{x \rightarrow x_0} f(x)$. If they are equal we can take the smallest δ and we can guarantee proof.

Left-hand, Right-hand, and Limits

Theorem

The $\lim_{x \rightarrow x_0} f(x)$ exists if and only if $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$

- Intuition that $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x) \Rightarrow \lim_{x \rightarrow x_0} f(x)$. If they are equal we can take the smallest δ and we can guarantee proof.
- Intuition that $\lim_{x \rightarrow x_0} f(x) \Rightarrow \lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$.
Absolute value is symmetric—so we must be converging from each side. (contradiction could work too!)

Left-hand, Right-hand, and Limits

Theorem

The $\lim_{x \rightarrow x_0} f(x)$ exists if and only if $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$

- Intuition that $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x) \Rightarrow \lim_{x \rightarrow x_0} f(x)$. If they are equal we can take the smallest δ and we can guarantee proof.
- Intuition that $\lim_{x \rightarrow x_0} f(x) \Rightarrow \lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$.
Absolute value is symmetric—so we must be converging from each side. (contradiction could work too!)
- We can also appeal to **sequences** to prove this stuff

Left-hand, Right-hand, and Limits

Theorem

The $\lim_{x \rightarrow x_0} f(x)$ exists if and only if $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$

- Intuition that $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x) \Rightarrow \lim_{x \rightarrow x_0} f(x)$. If they are equal we can take the smallest δ and we can guarantee proof.
- Intuition that $\lim_{x \rightarrow x_0} f(x) \Rightarrow \lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x)$.
Absolute value is symmetric—so we must be converging from each side. (contradiction could work too!)
- We can also appeal to **sequences** to prove this stuff

Trick: we'll show limits don't exist by showing

$$\lim_{x \rightarrow x_0^-} f(x) \neq \lim_{x \rightarrow x_0^+} f(x)$$

Finding Limits

Finding Limits

Student: Justin. what the hell with the δ 's and ϵ 's? What the hell am I going to use this for?

Finding Limits

Student: Justin. what the hell with the δ 's and ϵ 's? What the hell am I going to use this for?

Justin: Limits are used constantly in political science. And getting comfortable with this notation (by seeing it many times) is important

Finding Limits

Student: Justin. what the hell with the δ 's and ϵ 's? What the hell am I going to use this for?

Justin: Limits are used constantly in political science. And getting comfortable with this notation (by seeing it many times) is important

Student: fine. How am I going to find the limit? I can't do a $\delta - \epsilon$ proof yet.

Finding Limits

Student: Justin. what the hell with the δ 's and ϵ 's? What the hell am I going to use this for?

Justin: Limits are used constantly in political science. And getting comfortable with this notation (by seeing it many times) is important

Student: fine. How am I going to find the limit? I can't do a $\delta - \epsilon$ proof yet.

Justin: yes, those take time. For this class, **graphing** will be critical.

Algebra of Limits

Theorem

Suppose $f : \mathfrak{R} \rightarrow \mathfrak{R}$ and $g : \mathfrak{R} \rightarrow \mathfrak{R}$ with limits A and B at x_0 . Then,

$$i.) \lim_{x \rightarrow x_0} (f(x) + g(x)) = \lim_{x \rightarrow x_0} f(x) + \lim_{x \rightarrow x_0} g(x) = A + B$$

$$ii.) \lim_{x \rightarrow x_0} f(x)g(x) = \lim_{x \rightarrow x_0} f(x) \lim_{x \rightarrow x_0} g(x) = AB$$

Suppose $g(x) \neq 0$ for all $x \in \mathfrak{R}$ and $B \neq 0$ then $\frac{f(x)}{g(x)}$ has a limit at x_0 and


$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow x_0} f(x)}{\lim_{x \rightarrow x_0} g(x)} = \frac{A}{B}$$

Challenge Problems

Suppose $\lim_{x \rightarrow x_0} f(x) = a$. Find $\lim_{x \rightarrow x_0} \frac{f(x)^3 + f(x)^2}{f(x)}$

Continuity


$$f(x) = (x^2 - 1)/(x - 1)$$



- Limit exists at 1

Continuity


$$f(x) = (x^2 - 1)/(x - 1)$$



- Limit exists at 1
- But hole in function

Continuity

$$f(x) = (x^2 - 1)/(x - 1)$$



- Limit exists at 1
- But hole in function
- Fails the **pencil** test, **discontinuous** at 1

Continuity, Rigorous Definition


Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ and consider $x_0 \in \mathbb{R}$. We will say f is continuous at x_0 if for each $\epsilon > 0$ there is a $\delta > 0$ such that if,


$$\begin{aligned} |x - x_0| &< \delta \text{ for all } x \in \mathbb{R} \text{ then} \\ |f(x) - f(x_0)| &< \epsilon \end{aligned}$$

- Previously $f(x_0)$ was replaced with L .
- Now: $f(x)$ has to converge on itself at x_0 .
- Continuity is more restrictive than limit

Examples




Examples




Examples

$$f(x) = x^2$$



Examples



Continuity and Limits

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ with $x_0 \in \mathbb{R}$. Then f is continuous at x_0 if and only if f has a limit at x_0 and that $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

Proof.

(\Rightarrow). Suppose f is continuous at x_0 . This implies that for each $\epsilon > 0$ there is $\delta > 0$ such that $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$. This is the definition of a limit, with $L = f(x_0)$.

(\Leftarrow). Suppose f has a limit at x_0 and that limit is $f(x_0)$. This implies that for each $\epsilon > 0$ there is $\delta > 0$ such that $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$. But this is the definition of continuity. □


Algebra of Continuous Functions

Theorem

Suppose $f : \mathcal{R} \rightarrow \mathcal{R}$ and $g : \mathcal{R} \rightarrow \mathcal{R}$ are continuous at x_0 . Then,

- i.) $f(x) + g(x)$ is continuous at x_0
- ii.) $f(x)g(x)$ is continuous at x_0
- iii. if $g(x_0) \neq 0$, then $\frac{f(x)}{g(x)}$ is continuous at x_0


Use theorem about limits to prove continuous theorems.




How Functions Change

- **Derivatives**—Rates of change in functions
- Foundational across a lot of work in Poli Sci.
- A special **limit**
- Cover three broad ideas
 - Geometric interpretation/intuition
 - Formulas/Algebra derivatives
 - Famous theorems


Rates of Change in a Function




Rates of Change in a Function




Rates of Change in a Function




Rates of Change in a Function




Rates of Change in a Function




Rates of Change in a Function



Rates of Change in a Function




Rates of Change in a Function




- Rate of Change
↔ Return on
Vote Share/\$
Invested

Rates of Change in a Function



- Rate of Change
~> Return on
Vote Share/\$
Invested
- Instantaneous
rate of change ~>
Increase in vote
share in response
to infinitesimally
small increase in
spending

Rates of Change in a Function



- Rate of Change
~> Return on Vote Share/\$ Invested
- Instantaneous rate of change ~> Increase in vote share in response to infinitesimally small increase in spending
- **Limit**

Derivative Definition

Suppose $f : \mathcal{R} \rightarrow \mathcal{R}$.

Definition

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

Definition

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

$$R(x) = \frac{f(x) - f(x_0)}{x - x_0}$$

Definition

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

$$R(x) = \frac{f(x) - f(x_0)}{x - x_0}$$

- $R(x)$ defines the rate of change.

Definition

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

$$R(x) = \frac{f(x) - f(x_0)}{x - x_0}$$

- $R(x)$ defines the rate of change.
- A derivative will examine what happens with a small perturbation at x_0

Definition

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

$$R(x) = \frac{f(x) - f(x_0)}{x - x_0}$$

- $R(x)$ defines the rate of change.
- A derivative will examine what happens with a small perturbation at x_0

Definition

Let $f : \mathbb{R} \rightarrow \mathbb{R}$. If the limit

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

$$R(x) = \frac{f(x) - f(x_0)}{x - x_0}$$

- $R(x)$ defines the rate of change.
- A derivative will examine what happens with a small perturbation at x_0

Definition

Let $f : \mathbb{R} \rightarrow \mathbb{R}$. If the limit

$$\begin{aligned} \lim_{x \rightarrow x_0} R(x) &= \frac{f(x) - f(x_0)}{x - x_0} \\ &= f'(x_0) \end{aligned}$$

Derivative Definition

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$. Measure rate of change at a point x_0 with a function $R(x)$,

$$R(x) = \frac{f(x) - f(x_0)}{x - x_0}$$

- $R(x)$ defines the rate of change.
- A derivative will examine what happens with a small perturbation at x_0

Definition

Let $f : \mathbb{R} \rightarrow \mathbb{R}$. If the limit

$$\begin{aligned}\lim_{x \rightarrow x_0} R(x) &= \frac{f(x) - f(x_0)}{x - x_0} \\ &= f'(x_0)\end{aligned}$$

exists then we say that f is **differentiable** at x_0 . If $f'(x_0)$ exists for all $x \in \text{Domain}$, then we say that f is differentiable.

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\lim_{x \rightarrow 1} R(x) = \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1}$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1}\end{aligned}$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1\end{aligned}$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1 \\ &= 2\end{aligned}$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1 \\ &= 2\end{aligned}$$

- Suppose $f(x) = |x|$ and consider $x_0 = 0$. Then,

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1 \\ &= 2\end{aligned}$$

- Suppose $f(x) = |x|$ and consider $x_0 = 0$. Then,

$$\lim_{x \rightarrow 0} R(x) = \lim_{x \rightarrow 0} \frac{|x|}{x}$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1 \\ &= 2\end{aligned}$$

- Suppose $f(x) = |x|$ and consider $x_0 = 0$. Then,

$$\lim_{x \rightarrow 0} R(x) = \lim_{x \rightarrow 0} \frac{|x|}{x}$$

$$\lim_{x \rightarrow 0^-} R(x) = -1$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1 \\ &= 2\end{aligned}$$

- Suppose $f(x) = |x|$ and consider $x_0 = 0$. Then,

$$\lim_{x \rightarrow 0} R(x) = \lim_{x \rightarrow 0} \frac{|x|}{x}$$

$$\lim_{x \rightarrow 0^-} R(x) = -1, \text{ but } \lim_{x \rightarrow 0^+} R(x) = 1.$$

Derivative Examples

- Suppose $f(x) = x^2$ and consider $x_0 = 1$. Then,

$$\begin{aligned}\lim_{x \rightarrow 1} R(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{(x - 1)(x + 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x + 1 \\ &= 2\end{aligned}$$

- Suppose $f(x) = |x|$ and consider $x_0 = 0$. Then,

$$\lim_{x \rightarrow 0} R(x) = \lim_{x \rightarrow 0} \frac{|x|}{x}$$

$\lim_{x \rightarrow 0^-} R(x) = -1$, but $\lim_{x \rightarrow 0^+} R(x) = 1$. So, not differentiable at 0.

Continuity and Derivatives

- $f(x) = |x|$ is **continuous** but not differentiable. This is because the change is **too abrupt**.
- Suggests **differentiability is a stronger condition**

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Continuity and Derivatives

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

Continuity and Derivatives

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

Continuity and Derivatives

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

$$f(x) = \frac{f(x) - f(x_0)}{x - x_0}(x - x_0) + f(x_0)$$

Continuity and Derivatives

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

$$\begin{aligned} f(x) &= \frac{f(x) - f(x_0)}{x - x_0}(x - x_0) + f(x_0) \\ &= R(x)(x - x_0) + f(x_0) \end{aligned}$$

Continuity and Derivatives

Theorem

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

$$\begin{aligned} f(x) &= \frac{f(x) - f(x_0)}{x - x_0}(x - x_0) + f(x_0) \\ &= R(x)(x - x_0) + f(x_0) \end{aligned}$$

If $f(x)$ is continuous at x_0 then, $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

Continuity and Derivatives

Theorem

Let $f : \mathfrak{R} \rightarrow \mathfrak{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

$$\begin{aligned} f(x) &= \frac{f(x) - f(x_0)}{x - x_0}(x - x_0) + f(x_0) \\ &= R(x)(x - x_0) + f(x_0) \end{aligned}$$

If $f(x)$ is continuous at x_0 then, $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

$$\lim_{x \rightarrow x_0} f(x) = \lim_{x \rightarrow x_0} [R(x)(x - x_0) + f(x_0)]$$

Continuity and Derivatives

Theorem

Let $f : \mathfrak{R} \rightarrow \mathfrak{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

$$\begin{aligned} f(x) &= \frac{f(x) - f(x_0)}{x - x_0}(x - x_0) + f(x_0) \\ &= R(x)(x - x_0) + f(x_0) \end{aligned}$$

If $f(x)$ is continuous at x_0 then, $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

$$\begin{aligned} \lim_{x \rightarrow x_0} f(x) &= \lim_{x \rightarrow x_0} [R(x)(x - x_0) + f(x_0)] \\ &= \left(\lim_{x \rightarrow x_0} R(x) \right) \left(\lim_{x \rightarrow x_0} (x - x_0) \right) + \lim_{x \rightarrow x_0} f(x_0) \end{aligned}$$

Continuity and Derivatives

Theorem

Let $f : \mathfrak{R} \rightarrow \mathfrak{R}$ be differentiable at x_0 . Then f is continuous at x_0 .

Proof.

This proof is all in the setup. Realize that,

$$\begin{aligned} f(x) &= \frac{f(x) - f(x_0)}{x - x_0}(x - x_0) + f(x_0) \\ &= R(x)(x - x_0) + f(x_0) \end{aligned}$$

If $f(x)$ is continuous at x_0 then, $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

$$\begin{aligned} \lim_{x \rightarrow x_0} f(x) &= \lim_{x \rightarrow x_0} [R(x)(x - x_0) + f(x_0)] \\ &= \left(\lim_{x \rightarrow x_0} R(x) \right) \left(\lim_{x \rightarrow x_0} (x - x_0) \right) + \lim_{x \rightarrow x_0} f(x_0) \\ &= f'(x_0)0 + f(x_0) = f(x_0) \end{aligned}$$

What goes wrong?

Consider the following piecewise function:

$$\begin{aligned}f(x) &= x^2 \text{ for all } x \in \mathbb{R} \setminus 0 \\f(x) &= 1000 \text{ for } x = 0\end{aligned}$$

Consider derivative at 0. Then,

$$\begin{aligned}\lim_{x \rightarrow 0} R(x) &= \lim_{x \rightarrow 0} \frac{f(x) - 1000}{x - 0} \\&= \lim_{x \rightarrow 0} \frac{x^2}{x} - \lim_{x \rightarrow 0} \frac{1000}{x}\end{aligned}$$

$\lim_{x \rightarrow 0} \frac{1000}{x}$ diverges, so the limit doesn't exist.

Calculating Derivatives

- **Rarely** will we take limit to calculate derivative.
- Rather, rely on **rules** and properties of derivatives
- **Important:** do not forget core intuition

Strategy:

- Algebra theorems
- Some specific derivatives
- Work on problems

Some Derivative Rules

Suppose a is some constant, $f(x)$ and $g(x)$ are functions

Some Derivative Rules

Suppose a is some constant, $f(x)$ and $g(x)$ are functions

$$f(x) = x \quad ; \quad f'(x) = 1$$

Some Derivative Rules

Suppose a is some constant, $f(x)$ and $g(x)$ are functions

$$f(x) = x \quad ; \quad f'(x) = 1$$

$$f(x) = ax^k \quad ; \quad f'(x) = (a)(k)x^{k-1}$$

Some Derivative Rules

Suppose a is some constant, $f(x)$ and $g(x)$ are functions

$$f(x) = x \quad ; \quad f'(x) = 1$$

$$f(x) = ax^k \quad ; \quad f'(x) = (a)(k)x^{k-1}$$

$$f(x) = e^x \quad ; \quad f'(x) = e^x$$

Some Derivative Rules

Suppose a is some constant, $f(x)$ and $g(x)$ are functions

$$f(x) = x \quad ; \quad f'(x) = 1$$

$$f(x) = ax^k \quad ; \quad f'(x) = (a)(k)x^{k-1}$$

$$f(x) = e^x \quad ; \quad f'(x) = e^x$$

$$f(x) = \sin(x) \quad ; \quad f'(x) = \cos(x)$$

Some Derivative Rules

Suppose a is some constant, $f(x)$ and $g(x)$ are functions

$$f(x) = x \quad ; \quad f'(x) = 1$$

$$f(x) = ax^k \quad ; \quad f'(x) = (a)(k)x^{k-1}$$

$$f(x) = e^x \quad ; \quad f'(x) = e^x$$

$$f(x) = \sin(x) \quad ; \quad f'(x) = \cos(x)$$

$$f(x) = \cos(x) \quad ; \quad f'(x) = -\sin(x)$$

Algebra of Derivatives

Theorem

Suppose $f : \mathcal{R} \rightarrow \mathcal{R}$ and $g : \mathcal{R} \rightarrow \mathcal{R}$ and both are differentiable at $x_0 \in \mathcal{R}$.

Then,

Algebra of Derivatives

Theorem

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ and both are differentiable at $x_0 \in \mathbb{R}$.

Then,

i) *$h(x) = f(x) + g(x)$ is differentiable at x_0 and*

Algebra of Derivatives

Theorem

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ and both are differentiable at $x_0 \in \mathbb{R}$.

Then,

i) *$h(x) = f(x) + g(x)$ is differentiable at x_0 and*

$$h'(x_0) = f'(x_0) + g'(x_0)$$

Algebra of Derivatives

Theorem

*Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ and both are differentiable at $x_0 \in \mathbb{R}$.
Then,*

i) $h(x) = f(x) + g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0) + g'(x_0)$$

ii) $h(x) = f(x)g(x)$ is differentiable at x_0 and

Algebra of Derivatives

Theorem

Suppose $f : \mathcal{R} \rightarrow \mathcal{R}$ and $g : \mathcal{R} \rightarrow \mathcal{R}$ and both are differentiable at $x_0 \in \mathcal{R}$.
Then,

i) $h(x) = f(x) + g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0) + g'(x_0)$$

ii) $h(x) = f(x)g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0)g(x_0) + g'(x_0)f(x_0)$$

Algebra of Derivatives

Theorem

Suppose $f : \mathcal{R} \rightarrow \mathcal{R}$ and $g : \mathcal{R} \rightarrow \mathcal{R}$ and both are differentiable at $x_0 \in \mathcal{R}$.
Then,

i) $h(x) = f(x) + g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0) + g'(x_0)$$

ii) $h(x) = f(x)g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0)g(x_0) + g'(x_0)f(x_0)$$

iii) $h(x) = \frac{f(x)}{g(x)}$ with $g(x) \neq 0$ then,

Algebra of Derivatives

Theorem

Suppose $f : \mathcal{R} \rightarrow \mathcal{R}$ and $g : \mathcal{R} \rightarrow \mathcal{R}$ and both are differentiable at $x_0 \in \mathcal{R}$.
Then,

i) $h(x) = f(x) + g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0) + g'(x_0)$$

ii) $h(x) = f(x)g(x)$ is differentiable at x_0 and

$$h'(x_0) = f'(x_0)g(x_0) + g'(x_0)f(x_0)$$

iii) $h(x) = \frac{f(x)}{g(x)}$ with $g(x) \neq 0$ then,

$$h'(x_0) = \frac{f'(x_0)g(x_0) - g'(x_0)f(x_0)}{g(x_0)^2}$$

Challenge Problems

Differentiate the following functions and evaluate at the specified value

1) $f(x) = x^3 + 5x^2 + 4x$, at $x_0 = 2$

2) $f(x) = \sin(x)x^3$ at $x_0 = y$

3) $f(x) = \frac{e^x}{x^3}$ at $x = 2$

4) $g(x) = \log(x)x^3$ at $x = x_0$

5) Suppose $f(x) = x^2$ and $g(x) = x^3$. Find all x such that $f'(x) > g'(x)$.

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Suppose $k > 0$. We will proceed by induction. Suppose $k = 1$, $f(x) = x$

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Suppose $k > 0$. We will proceed by induction. Suppose $k = 1$, $f(x) = x$

$$f'(\tilde{x}) = \lim_{x \rightarrow \tilde{x}} \frac{x - \tilde{x}}{x - \tilde{x}}$$

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Suppose $k > 0$. We will proceed by induction. Suppose $k = 1$, $f(x) = x$

$$\begin{aligned} f'(\tilde{x}) &= \lim_{x \rightarrow \tilde{x}} \frac{x - \tilde{x}}{x - \tilde{x}} \\ &= 1 = 1x^0 \end{aligned}$$

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Suppose $k > 0$. We will proceed by induction. Suppose $k = 1$, $f(x) = x$

$$\begin{aligned} f'(\tilde{x}) &= \lim_{x \rightarrow \tilde{x}} \frac{x - \tilde{x}}{x - \tilde{x}} \\ &= 1 = 1x^0 \end{aligned}$$

Suppose theorem holds for $k = r$, $f(x) = x^r$. Consider $g(x) = x^{r+1}$. We know that $g(x) = f(x)x$. By product rule,

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Suppose $k > 0$. We will proceed by induction. Suppose $k = 1$, $f(x) = x$

$$\begin{aligned} f'(\tilde{x}) &= \lim_{x \rightarrow \tilde{x}} \frac{x - \tilde{x}}{x - \tilde{x}} \\ &= 1 = 1x^0 \end{aligned}$$

Suppose theorem holds for $k = r$, $f(x) = x^r$. Consider $g(x) = x^{r+1}$. We know that $g(x) = f(x)x$. By product rule,

$$g'(x) = f(x)x' + f'(x)x = x^r 1 + rx^{r-1}x$$

Proving Property of Derivatives

Theorem

Suppose $f(x) = x^k$ and k is a positive integer. If $k = 0$ then $f'(x) = 0$. If $k > 0$, then, $f'(x) = kx^{k-1}$.

Proof.

If $k = 0$ then, $x^k = 1$. The $\lim_{x \rightarrow \tilde{x}} \frac{1-1}{x-\tilde{x}} = 0$.

Suppose $k > 0$. We will proceed by induction. Suppose $k = 1$, $f(x) = x$

$$\begin{aligned} f'(\tilde{x}) &= \lim_{x \rightarrow \tilde{x}} \frac{x - \tilde{x}}{x - \tilde{x}} \\ &= 1 = 1x^0 \end{aligned}$$

Suppose theorem holds for $k = r$, $f(x) = x^r$. Consider $g(x) = x^{r+1}$. We know that $g(x) = f(x)x$. By product rule,

$$\begin{aligned} g'(x) = f(x)x' + f'(x)x &= x^r 1 + rx^{r-1}x \\ &= x^r + rx^r = (r+1)x^r \end{aligned}$$

Chain Rule

Common to have functions in functions

$$\begin{aligned} f(x) &= \frac{e^{-\frac{(x-\mu)^2}{2\sigma^2}}}{\sqrt{2\pi}} \\ &= \frac{f(g(x))}{\sqrt{2\pi}} \end{aligned}$$

To deal with this, we use the **chain rule**

Theorem

Suppose $g : \mathfrak{R} \rightarrow \mathfrak{R}$ and $f : \mathfrak{R} \rightarrow \mathfrak{R}$. Suppose both $f(x)$ and $g(x)$ are differentiable at x_0 . Define $h(x) = g(f(x))$. Then,

$$h'(x_0) = g'(f(x_0))f'(x_0)$$

Examples of Chain Rule in Action

- $h(x) = e^{2x}$. $g(x) = e^x$. $f(x) = 2x$. So
 $h(x) = g(f(x)) = g(2x) = e^{2x}$. Taking derivatives, we have

$$h'(x) = g'(f(x))f'(x) = e^{2x}2$$

- $h(x) = \log(\cos(x))$. $g(x) = \log(x)$. $f(x) = \cos(x)$.
 $h(x) = g(f(x)) = g(\cos(x)) = \log(\cos(x))$

$$h'(x) = g'(f(x))f'(x) = \frac{-1}{\cos(x)} \sin(x) = -\tan(x)$$

Derivatives and Properties of Functions

Derivatives reveal an **immense** amount about functions

- Often use to **optimize** a function (tomorrow)
- But also reveal **average rates of change**
- Or crucial properties of functions

Goal: introduce ideas. Hopefully make them less shocking when you see them in work


Relative Maxima, Minima and Derivatives

Theorem

Suppose $f : [a, b] \rightarrow \mathfrak{R}$. Suppose f has a relative maxima or minima on (a, b) and call that $c \in (a, b)$. Then $f'(c) = 0$.

Intuition:

Rolle's Theorem



Relative Maxima, Minima and Derivatives

Theorem

Rolle's Theorem Suppose $f : [a, b] \rightarrow \mathfrak{R}$ and f is continuous on $[a, b]$ and differentiable on (a, b) . Then if $f(a) = f(b) = 0$, there is $c \in (a, b)$ such that $f'(c) = 0$.

Proof **Intuition** Consider (WLOG) a relative maximum c . Consider the left-hand and right-hand limits

$$\lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \geq 0$$
$$\lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \leq 0$$

Theorem

Rolle's Theorem Suppose $f : [a, b] \rightarrow \mathfrak{R}$ and f is continuous on $[a, b]$ and differentiable on (a, b) . Then if $f(a) = f(b) = 0$, there is $c \in (a, b)$ such that $f'(c) = 0$.

But we also know that

$$\lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} = f'(c)$$
$$\lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} = f'(c)$$

The only way, then, that

$$\lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} = \lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \text{ is if } f'(c) = 0.$$

What Goes Up Must Come Down

Theorem


Rolle's Theorem Suppose $f : [a, b] \rightarrow \mathfrak{R}$ and f is continuous on $[a, b]$ and differentiable on (a, b) . Then if $f(a) = f(b) = 0$, there is $c \in (a, b)$ such that $f'(c) = 0$.

What Goes Up Must Come Down

Theorem

Rolle's Theorem Suppose $f : [a, b] \rightarrow \mathfrak{R}$ and f is continuous on $[a, b]$ and differentiable on (a, b) . Then if $f(a) = f(b) = 0$, there is $c \in (a, b)$ such that $f'(c) = 0$.

Rolle's Theorem



Mean Value Theorem


Theorem

If $f : [a, b] \rightarrow \mathfrak{R}$ is continuous on $[a, b]$ and differentiable on (a, b) , then there is a $c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$


Rolle's Theorem, Rotated

Rolle's Theorem



Rolle's Theorem, Rotated

Mean Value Theorem



Why You Should Care

- 1) This will come up in a formal theory article. You'll at least know where to look
- 2) It allows us to say lots of powerful stuff about functions

Powerful Applications of Mean Value Theorem

Theorem

Suppose that $f : [a, b] \rightarrow \mathfrak{R}$ is continuous on $[a, b]$ and differentiable on (a, b) . Then,

- i) If $f'(x) \neq 0$ for all $x \in (a, b)$ then f is 1-1
- ii) If $f'(x) = 0$ then $f(x)$ is constant
- iii) If $f'(x) > 0$ for all $x \in (a, b)$ then f is strictly increasing
- iv) If $f'(x) < 0$ for all $x \in (a, b)$ then f is strictly decreasing

Let's prove these in turn

- Why—because they are just about applying ideas

If $f'(x) \neq 0$ for all $x \in (a, b)$ then f is 1-1

By way of contradiction, suppose that f is not 1-1. Then there is $x, y \in (a, b)$ such that $f(x) = f(y)$. Then,

$$f'(c) = \frac{f(x) - f(y)}{x - y} = \frac{0}{x - y} = 0$$

If $f'(x) \neq 0$ for all $x \in (a, b)$ then f is 1-1



If $f'(x) \neq 0$ for all $x \in (a, b)$ then f is 1-1



$f' \neq 0$ for all x !

If $f'(x) = 0$ then $f(x)$ is constant

By way of contradiction, suppose that there is $x, y \in (a, b)$ such that $f(x) \neq f(y)$. But then,

$$f'(c) = \frac{f(x) - f(y)}{x - y} \neq 0$$

contradiction

If $f'(x) > 0$ for all $x \in (a, b)$ then then f is strictly increasing

By way of contradiction, suppose that there is $x, y \in (a, b)$ with $y < x$ but $f(y) > f(x)$. But then,

$$f'(c) = \frac{f(x) - f(y)}{x - y} < 0$$

contradiction

Bonus: proof for strictly decreasing

Approximating functions and second order conditions

Theorem

Taylor's Theorem Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x)$ is infinitely differentiable function. Then, the Taylor expansion of $f(x)$ around a is given by

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$

$$f(x) = \sum_{n=0}^{\infty} \frac{f^n(a)}{n!}(x-a)^n$$

Example Function

Suppose $a = 0$ and $f(x) = e^x$. Then,

$$f'(x) = e^x$$

$$f''(x) = e^x$$

$$\vdots \quad \vdots \quad \vdots$$

$$f^n(x) = e^x$$

This implies

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} \dots + \frac{x^n}{n!} + \dots$$

Wrap up

Lots of territory.

What are your questions?

This Week

Lab Tonight!