

# Vectors in Julia

ENGR108  
Stanford University

April 3, 2023

# Vectors in Julia

main topics:

- ▶ how to create and manipulate vectors in Julia
- ▶ how Julia notation differs from math notation

# Outline

Vectors

Vector operations

Norm and distance

## Vectors

- ▶ vectors are represented by arrays in Julia
- ▶ to create the 3-vector

$$x = (8, -4, 3.5) = \begin{bmatrix} 8 \\ -4 \\ 3.5 \end{bmatrix}$$

use

`x = [8, -4, 3.5]`

(`x = [8;-4;3.5]` also works)

- ▶ watch out for similar looking expressions
  - `(8,-4,3.5)` and `[8 -4 3.5]` are not equivalent in Julia
- ▶ length of a vector: `length(x)`

## Ranges

- ▶ to get a range from  $i$  to  $j$  (for  $i \leq j$ ), use a colon (`:`)
  - the range from 1 to 10 is `1:10`
  - `collect(1:10)` returns the array
- ▶ the default increment between values is 1. (`1:3` is 1, 2, 3)
- ▶ to specify an increment size add an additional argument:
  - the range from 1 to 10 with a step size of 0.1 is `1:0.1:10`

## Indexing and slicing

- ▶ indexes run from 1 to  $n$ :  $x_2$  is `x[2]`
- ▶ can also set an element, e.g., `x[3] = 10.5`
- ▶ use a range to select more than one element
  - `x[2:3]` selects the second and third elements
- ▶ `x[end]` selects the last element
- ▶ to select every other element use `x[1:2:end]`

## Block vectors

- ▶ to form a stacked vector like

$$a = (b, c) = \begin{bmatrix} b \\ c \end{bmatrix}$$

(with  $b$  and  $c$  vectors)

$$a = [b; c]$$

( $a = [b, c]$  does NOT work)

- ▶ can mix vectors and scalars:

$$a = [b; 2; c; -6]$$

## Basic functions for arrays

- ▶ sum of (the entries of) a vector: `sum(x)`
- ▶ mean of the entries (`avg(x)`): `mean(x)`
- ▶  $0_n$  is `zeros(n)`
- ▶  $\mathbf{1}_n$  is `ones(n)`



## Creating unit vectors

- ▶ form  $e_3$  with length 10
- ▶ create a zero vector of size 10 then set the third element to 1  
`e_3 = zeros(10); e_3[3] = 1;`

## List of vectors

- ▶ to form a list with vectors a, b, and c:  
`vector_list = [a,b,c]`
- ▶ the second vector in this list is `vector_list[2]`
- ▶ to access an element in a vector: `vector_list[2][3]`

# Outline

Vectors

Vector operations

Norm and distance

## Vector addition and subtraction

- ▶ vector addition uses +, for example

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$$

is written

$$[1, 2, 3] + [4, 5, 6]$$

- ▶ subtraction uses -
- ▶ the arrays must have the same length (unless one is scalar)

## Scalar-vector addition

- ▶ in Julia, a scalar and a vector can be added
- ▶ the scalar is added to each entry of the vector

`[2, 4, 8] + 3`

gives (in mathematical notation)

$$\begin{bmatrix} 2 \\ 4 \\ 8 \end{bmatrix} + 3\mathbf{1} = \begin{bmatrix} 5 \\ 7 \\ 11 \end{bmatrix}$$

## Scalar-vector multiplication

- ▶ scalar-vector multiplication uses \*
- ▶ for example,

$$(-2) \begin{bmatrix} 1 \\ 9 \\ 6 \end{bmatrix}$$

is written

$$-2 * [1, 9, 6]$$

- ▶ the other order gives the same result:  
 $[1, 9, 6] * -2$

## Inner product

- ▶ inner product  $a^T b$  is written as  $\text{dot}(a, b)$
- ▶  $a$  and  $b$  must have the same length

# Outline

Vectors

Vector operations

Norm and distance



## Norm and distance

- ▶ the norm  $\|x\| = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}$  is written  
norm(x)
- ▶ **dist**( $x, y$ ) =  $\|x - y\|$  is written  
norm(x-y)

## RMS value

- ▶  $\mathbf{rms}(x)$  is defined as

$$\mathbf{rms}(x) = \sqrt{\frac{1}{n} (x_1^2 + \dots + x_n^2)} = \frac{\|x\|}{\sqrt{n}}.$$

- ▶ can be expressed as

$$\mathbf{rms\_x} = \mathbf{norm}(x)/\mathbf{sqrt}(\mathbf{length}(x))$$

## Standard deviation

- ▶ standard deviation is defined as

$$\mathbf{std}(x) = \frac{\|x - \mathbf{avg}(x)\mathbf{1}\|}{\sqrt{n}}$$

- ▶ which can be expressed as

$$\mathbf{std\_of\_x} = \mathbf{norm}(x - \mathbf{mean}(x))/\mathbf{sqrt}(\mathbf{length}(x))$$

- ▶ warning: the Julia function `std` does *not* use this definition

# Angle

- ▶ the angle between two vectors  $a$  and  $b$  is

$$\angle(a, b) = \arccos\left(\frac{a^T b}{\|a\| \|b\|}\right)$$

- ▶ can be expressed as

$$\text{angle\_a\_b} = \text{acos}(\text{dot}(a, b) / (\text{norm}(a) * \text{norm}(b)))$$